

THE SEMILATTICE, BOX, AND LATTICE-TENSOR PRODUCTS IN QUANTUM LOGIC

BORIS ISCHI

ABSTRACT. Given two complete atomistic lattices $\mathcal{L}_1$ and $\mathcal{L}_2$, we define a set $\mathbf{S} = \mathbf{S}(\mathcal{L}_1, \mathcal{L}_2)$ of complete atomistic lattices by means of three axioms (natural regarding the description of *separated* quantum compound systems), or in terms of a universal property with respect to a given class of bimorphisms. We prove that $\mathbf{S}$ is a complete lattice. The bottom element $\mathcal{L}_1 \mathbin{\bigtriangleup} \mathcal{L}_2$ is the *separated product* of Aerts. For atomistic lattices with 1 (not complete), $\mathcal{L}_1 \mathbin{\bigtriangleup} \mathcal{L}_2 \cong \mathcal{L}_1 \square \mathcal{L}_2$ the *box product* of Grätzer and Wehrung, and, in case $\mathcal{L}_1$ and $\mathcal{L}_2$ are moreover coatomistic, $\mathcal{L}_1 \mathbin{\bigtriangleup} \mathcal{L}_2 \cong \mathcal{L}_1 \boxtimes \mathcal{L}_2$ the *lattice tensor product*. The top element $\mathcal{L}_1 \mathbin{\bigtriangledown} \mathcal{L}_2$ is the (complete) join-semilattice tensor product of Fraser, which is isomorphic to the tensor products of Chu and Shmueli. With some additional hypotheses on $\mathcal{L}_1$ and $\mathcal{L}_2$ (true if $\mathcal{L}_1$ and $\mathcal{L}_2$ are moreover orthomodular with the covering property), we prove that $\mathbf{S}$ is a singleton if and only if $\mathcal{L}_1$ or $\mathcal{L}_2$ is distributive, if and only if $\mathcal{L}_1 \mathbin{\bigtriangledown} \mathcal{L}_2$ has the covering property. Our main result reads: $\mathcal{L} \in \mathbf{S}$ admits an orthocomplementation if and only if $\mathcal{L} = \mathcal{L}_1 \mathbin{\bigtriangleup} \mathcal{L}_2$. For $\mathcal{L}_1$ and $\mathcal{L}_2$ moreover irreducible, we characterize the automorphisms of each $\mathcal{L} \in \mathbf{S}$ in terms of those of $\mathcal{L}_1$ and $\mathcal{L}_2$. At the end, we construct an example $\mathcal{L}_1 \mathbin{\bigoplus} \mathcal{L}_2$ in $\mathbf{S}$ which has the covering property.

1. INTRODUCTION

In quantum logic, one associates to a physical system S a couple $(\Sigma_S, \mathcal{L}_S \subseteq 2^{\Sigma_S})$, where Σ_S represents all possible states of S and $\mathcal{L}_S$ the experimental propositions concerning S : A proposition represented by some $a \in \mathcal{L}_S$ is true with probability 1 if and only if the state of S lies in a [4].

A cornerstone is the following theorem, by which the Hilbert space structure of quantum mechanics can be recovered from a certain number of axioms on the poset $\mathcal{L}_S$ [15] (see [13], Theorems 34.5 and 34.9).

Theorem 1.1. *If $\mathcal{L}_S$ is an irreducible orthocomplemented simple closure space on Σ_S (i.e. $\mathcal{L}_S$ is closed under arbitrary set-intersections and contains $\emptyset$, Σ_S and all singletons) with the covering property of length ≥ 4 , then there is a $*$ -division ring $\mathbb{K}$ and a vector space E_S over $\mathbb{K}$ with an Hermitian form such that $\mathcal{L}_S$ is orthoisomorphic to the lattice $\mathbf{P}(E_S)$ of E_S -closed subspaces of E_S . Moreover, if $\mathbb{K} = \mathbb{R}$ or $\mathbb{C}$ with the usual involution, then E_S is a Hilbert space if and only if $\mathcal{L}_S$ is also orthomodular.*

If S is a compound system consisting of two subsystems S_1 and S_2 , then $\Sigma_{S_1} \times \Sigma_{S_2} \subseteq \Sigma_S$. In classical mechanics there is equality whereas in quantum theory Σ_S

1991 *Mathematics Subject Classification.* Primary 06B23, 06C15; Secondary 81P10.

Key words and phrases. Complete atomistic ortholattice, tensor product, quantum logic.

Supported by the Swiss National Science Foundation.

is given by all one-dimensional subspaces of the Hilbert space $\mathcal{H}_S = \mathcal{H}_{S_1} \otimes \mathcal{H}_{S_2}$ [14]. However, when two quantum systems S_1 and S_2 are *separated* we have [1, 11]

$$(P1) \quad \Sigma_S = \Sigma_{S_1} \times \Sigma_{S_2}.$$

In particular, simultaneous experiments on both systems can be performed and any experiment done on one system does not alter the state of the other system. Therefore, if P_1 is a proposition concerning S_1 represented by some $a_1 \in \mathcal{L}_1$ and P_2 is a proposition concerning S_2 represented by some $a_2 \in \mathcal{L}_2$, then the proposition $P_1 \text{ OR } P_2$ concerning the compound system is true with probability 1 if and only if the state p_1 of S_1 lies in a_1 or the state p_2 of S_2 lies in a_2 . In other words [1, 11]

$$(P2) \quad a_1 \times \Sigma_{S_2} \cup \Sigma_{S_1} \times a_2 \in \mathcal{L}_S, \text{ for all } a_1 \in \mathcal{L}_{S_1} \text{ and } a_2 \in \mathcal{L}_{S_2}.$$

For separated systems, it can be shown that $\mathcal{L}_S$, which we denote by $\mathcal{L}_{S_{sep}}$, cannot be isomorphic to the lattice of closed subspaces of a Hilbert space [2]. As consequence some axioms of Theorem 1.1 are failing in $\mathcal{L}_{S_{sep}}$.

In [1], Aerts proposed a model $\mathcal{L}_{S_1} \oslash \mathcal{L}_{S_2}$ for $\mathcal{L}_{S_{sep}}$, call the *separated product*. For $\mathcal{L}_{S_1}$ and $\mathcal{L}_{S_2}$ orthocomplemented simple closure spaces (in that case the separated product is an orthocomplemented simple closure space), he proved that if $\mathcal{L}_{S_1} \oslash \mathcal{L}_{S_2}$ has the covering property or is orthomodular, then $\mathcal{L}_{S_1}$ or $\mathcal{L}_{S_2}$ is a power set, hence S_1 or S_2 is a classical system [1]. As a consequence, according to Aerts, the covering property and orthomodularity do not hold in $\mathcal{L}_{S_{sep}}$. A similar conclusion concerning orthomodularity was obtained by Pulmannová in [17].

Here we show that $\mathcal{L}_{S_{sep}}$ does not admit an orthocomplementation. Moreover we provide a natural model for $\mathcal{L}_{S_{sep}}$ which has the covering property. We proceed as follows: Following [16] and [1], we assume that $\mathcal{L}_{S_1}$, $\mathcal{L}_{S_2}$ and $\mathcal{L}_{S_{sep}}$ are simple closure spaces. Moreover, in addition to Axioms P1 and P2 above, we postulate that

$$(P3) \quad \text{for all } p_i \in \Sigma_{S_i} \text{ and all } A_i \subseteq \Sigma_{S_i}, [p_1 \times A_2 \in \mathcal{L}_{S_{sep}} \Rightarrow A_2 \in \mathcal{L}_2] \text{ and } [A_1 \times p_2 \in \mathcal{L}_{S_{sep}} \Rightarrow A_1 \in \mathcal{L}_1].$$

We refer to [11] for detailed physical justifications of Axioms P1-P3. Let $S_{12} = S(\mathcal{L}_{S_1}, \mathcal{L}_{S_2})$ be the set of all simple closure spaces on $\Sigma_{S_1} \times \Sigma_{S_2}$ for which Axioms P2 and P3 hold. Let $\mathcal{L} \in S_{12}$. We prove that if $\mathcal{L}_{S_1}$ and $\mathcal{L}_{S_2}$ are orthocomplemented simple closure spaces with the covering property, then $\mathcal{L}$ admits an orthocomplementation if and only if $\mathcal{L} = \mathcal{L}_{S_1} \oslash \mathcal{L}_{S_2}$. We conclude by a simple physical argument which shows that certainly $\mathcal{L}_{S_1} \oslash \mathcal{L}_{S_2} \subsetneq \mathcal{L}_{S_{sep}}$ [11].

The rest of the paper is organized as follows. In Section 2 we fix some basic terminology and notations and we define $S \equiv S(\mathcal{L}_1, \dots, \mathcal{L}_n)$. We prove that S is a complete lattice, the bottom and top elements of which are denoted by $\bigwedge_i \mathcal{L}_i$ and $\bigvee_i \mathcal{L}_i$ respectively. In Section 3 we compare $\oslash$ and $\bigvee$ to a certain number of tensor products. An equivalent definition of S , in terms of a universal property with respect to a given class of bimorphisms (arbitrary joins are preserved) is given in Section 4. Let $\mathcal{L} \in S$. In Section 6 we prove some basic relations between the central elements of $\mathcal{L}_i$'s and $\mathcal{L}$.

Further, with some additional hypotheses on each $\mathcal{L}_i$ (true if $\mathcal{L}_i$ is moreover orthomodular with the covering property) we prove that S is a singleton (*i.e.* $\bigwedge_i \mathcal{L}_i = \bigvee_i \mathcal{L}_i$) if and only if at most one $\mathcal{L}_i$ is not a power set (Section 5) if and only if $\bigwedge_i \mathcal{L}_i$ or $\bigvee_i \mathcal{L}_i$ has the covering property (Section 9). Recall that in the physical interpretation, $\mathcal{L}_{S_i} = 2^{\Sigma_{S_i}}$ if and only if S_i is classical.

Let $\mathcal{L} \in \mathbf{S}$ and $u : \mathcal{L} \rightarrow \mathcal{L}$ preserving arbitrary joins and sending atoms to atoms. In Section 7 we prove, under some hypotheses on the image of u , that if each $\mathcal{L}_i$ is moreover irreducible, then there is a permutation f and join-preserving maps $v_i : \mathcal{L}_i \rightarrow \mathcal{L}_{f(i)}$ sending atoms to atoms, such that on atoms $u = f \circ (v_1 \times \cdots \times v_n)$. A time evolution can be modelled by a map preserving arbitrary joins and sending atoms to atoms [6]. Hence, in the physical interpretation, this result shows that separated quantum systems remain separated only if they do not interact. Finally, Section 8 is devoted to our main result and Section 10 to the example mentioned above.

2. MAIN DEFINITIONS

In this section we give our main definitions. We start with some background material and basic notations used in the sequel.

Definition 2.1. By a *simple closure space* $\mathcal{L}$ (on Σ) we mean a set of subsets of a nonempty set Σ , ordered by set-inclusion, closed under arbitrary set-intersections (i.e. for all $\omega \subseteq \mathcal{L}$, $\cap \omega \in \mathcal{L}$), and containing Σ , $\emptyset$, and all singletons. We denote the bottom ($\emptyset$) and top (Σ) elements by 0 and 1 respectively. For $p \in \Sigma$, we denote the singleton $\{p\}$ by p . Hence $p \cup q$ stands for $\{p, q\}$.

Remark 2.2. Let $\mathcal{L}$ be a simple closure space on a (nonempty) set Σ . Then $\mathcal{L}$ is a complete atomistic lattice, the atoms of which correspond to all singletons of Σ . Note that if $A \subseteq \Sigma$, then $\vee_{\mathcal{L}}(A) = \cap\{b \in \mathcal{L} ; A \subseteq b\}$.

Conversely, let $\mathcal{L}$ be a complete atomistic lattice. Let Σ denote the set of atoms of $\mathcal{L}$, and, for each $a \in \mathcal{L}$, let $\Sigma[a]$ denote the set of atoms under a . Then $\{\Sigma[a] ; a \in \mathcal{L}\}$ is a simple closure space on Σ , isomorphic to $\mathcal{L}$.

Definition 2.3. A lattice $\mathcal{L}$ with 0 and 1 is *orthocomplemented* if there is a unary operation $^\perp$ (orthocomplementation), also denoted by $'$, such that for all $a, b \in \mathcal{L}$, $(a^\perp)^\perp = a$, $a \leq b$ implies $b^\perp \leq a^\perp$, and $a \vee a^\perp = 1$.

An orthocomplemented lattice is *orthomodular* if for all $a, b \in \mathcal{L}$, $a \leq b$ implies $b = a \vee (b \wedge a^\perp)$.

A lattice with 0 has the *covering property* if for any atom p and any $a \in \mathcal{L}$, $p \wedge a = 0$ implies that $p \vee a$ covers a (in symbols $p \vee a \succ a$).

Finally, a lattice $\mathcal{L}$ with 0 and 1 is called a *DAC-lattice* if $\mathcal{L}$ and its dual $\mathcal{L}^*$ (defined by the converse order-relation) are atomistic with the covering property [13]. We say that a lattice $\mathcal{L}$ with 1 is *coatomistic* if the dual $\mathcal{L}^*$ is atomistic.

Remark 2.4. Note that an orthocomplemented atomistic lattice with the covering property is a DAC-lattice. Note also that in Theorem 1.1, if instead of orthocomplemented, the simple closure space is a DAC-lattice, then there is a pair of dual vector spaces such that a representation theorem similar to Theorem 1.1 holds (see [13], Theorem 33.7).

Definition 2.5. We denote the category of simple closure spaces with maps preserving arbitrary joins (hence 0) by $\mathbf{Cl}$, and the sub-category of simple closure spaces on a particular (nonempty) set Σ , by $\mathbf{Cl}(\Sigma)$. $\mathbf{2}$ denotes the simple closure space isomorphic to the two-element lattice.

Let $\mathcal{L} \in \mathbf{Cl}(\Sigma)$. We write $\text{Aut}(\mathcal{L})$ for the group of automorphisms of $\mathcal{L}$. Let $\mathcal{L}_1 \in \mathbf{Cl}(\Sigma_1)$. Note that any map $u : \mathcal{L} \rightarrow \mathcal{L}_1$ sending atoms to atoms induces

a mapping from Σ to Σ_1 , which we also call u . Thus, if $u \in \text{Aut}(\mathcal{L})$, then for all $a \in \mathcal{L}$, $u(a) = \{u(p); p \in a\}$.

If $\mathcal{L}$ is orthocomplemented, for $p, q \in \Sigma$, we write $p \perp q$ if and only if $p \in q^\perp$, where $q^\perp$ stands for $\{q\}^\perp$.

Finally, if $\mathcal{H}$ is a complex Hilbert space, then $\Sigma_{\mathcal{H}}$ denotes the set of one-dimensional subspaces of $\mathcal{H}$ and $\mathbf{P}(\mathcal{H})$ stands for the simple closure space isomorphic to the lattice of closed subspaces of $\mathcal{H}$. Moreover, we write $\mathbf{U}(\mathcal{H})$ for the set of automorphisms of $\mathbf{P}(\mathcal{H})$ induced by unitary maps on $\mathcal{H}$.

Definition 2.6. Let $\{\Sigma_\alpha\}_{\alpha \in \Omega}$ be a family of nonempty sets, $\Sigma = \prod_\alpha \Sigma_\alpha$, $\beta \in \Omega$, $p \in \Sigma$, and $R \subseteq \Sigma$. We shall make use of the following notations:

- (1) We denote by $\pi_\beta : \Sigma \rightarrow \Sigma_\beta$ the β -th coordinate map, *i.e.*, $\pi_\beta(p) = p_\beta$.
- (2) We denote by $p[-, \beta] : \Sigma_\beta \rightarrow \Sigma$ the map which sends $q \in \Sigma_\beta$ to the element of Σ obtained by replacing p 's β -th entry by q .
- (3) We define $R_\beta[p] = \pi_\beta(p[\Sigma_\beta, \beta] \cap R)$.

We omit the β in $p[-, \beta]$ when no confusion can occur. For instance, we write $p[\Sigma_\beta]$ instead of $p[\Sigma_\beta, \beta]$.

Remark 2.7. $p[R_\beta[p]] = p[\Sigma_\beta] \cap R$.

Definition 2.8. Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$. We denote by $\mathbf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$ the set all $\mathcal{L} \in \mathbf{Cl}(\Sigma)$ such that

- (P1) $\Sigma = \prod_\alpha \Sigma_\alpha$,
- (P2) $\cup_\alpha \pi_\alpha^{-1}(a_\alpha) \in \mathcal{L}$, for all $a \in \prod_\alpha \mathcal{L}_\alpha$,
- (P3) for all $p \in \Sigma$, $\beta \in \Omega$, and $A \subseteq \Sigma_\beta$, $[p[A, \beta]] \in \mathcal{L} \Rightarrow A \in \mathcal{L}_\beta$.

Let $T = \prod_\alpha T_\alpha$ with $T_\alpha \subseteq \text{Aut}(\mathcal{L}_\alpha)$. We denote by $\mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$ the set of all $\mathcal{L} \in \mathbf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$ such that

- (P4) for all $v \in T$, there is $u \in \text{Aut}(\mathcal{L})$ such that $u(p)_\alpha = v_\alpha(p_\alpha)$ for all $p \in \Sigma$ and $\alpha \in \Omega$.

We call elements of $\mathbf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$ *weak tensor products*.

Definition 2.9. We say that $\mathcal{L} \in \mathbf{Cl}(\Sigma)$ (respectively $T \subseteq \text{Aut}(\mathcal{L})$) is *transitive* if the action of $\text{Aut}(\mathcal{L})$ (respectively the action of T) on Σ is transitive.

Remark 2.10. The u in Axiom P4 is necessarily unique. If all T_α in Axiom P4 are transitive, then $\mathcal{L}$ is transitive. Note also that, $\mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega) \subseteq \mathbf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$ for all $T = \prod_\alpha T_\alpha$, with $T_\alpha \subseteq \text{Aut}(\mathcal{L}_\alpha)$. The name “weak tensor product” is justified by the fact that $\mathbf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$ and $\mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$ can be defined in terms of a universal property with respect to a given class of bimorphisms of $\mathbf{Cl}$ (see Section 4).

Definition 2.11. Let $\{\Sigma_\alpha\}_{\alpha \in \Omega}$ be a family of nonempty sets and $\{\mathcal{L}_\alpha \subseteq 2^{\Sigma_\alpha}\}_{\alpha \in \Omega}$. Let $\Sigma = \prod_\alpha \Sigma_\alpha$. Then

$$\begin{aligned} \bigotimes \{\mathcal{L}_\alpha; \alpha \in \Omega\} &:= \{\cap \omega; \omega \subseteq \{\cup_\alpha \pi_\alpha^{-1}(a_\alpha); a \in \prod_\alpha \mathcal{L}_\alpha\}\}, \\ \bigvee \{\mathcal{L}_\alpha; \alpha \in \Omega\} &:= \{R \subseteq \Sigma; R_\beta[p] \in \mathcal{L}_\beta, \forall p \in \Sigma, \beta \in \Omega\}, \end{aligned}$$

ordered by set-inclusion.

Proposition 2.12. *Let $\{\Sigma_\alpha\}_{\alpha \in \Omega}$ be a family of nonempty sets, and let $\{\mathcal{L}_\alpha \subseteq 2^{\Sigma_\alpha}\}_{\alpha \in \Omega}$. Let $\{\Omega_\gamma \subseteq \Omega; \gamma \in \Gamma\}$ such that $\Omega = \coprod\{\Omega_\gamma; \gamma \in \Gamma\}$. Then*

$$\begin{aligned}\bigotimes\{\mathcal{L}_\alpha; \alpha \in \Omega\} &= \bigotimes_{\gamma \in \Gamma}(\bigotimes_{\alpha \in \Omega_\gamma} \mathcal{L}_\alpha), \\ \bigvee\{\mathcal{L}_\alpha; \alpha \in \Omega\} &= \bigvee_{\gamma \in \Gamma}(\bigvee_{\alpha \in \Omega_\gamma} \mathcal{L}_\alpha).\end{aligned}$$

Proof. Direct from Definition 2.11. $\square$

Theorem 2.13. *Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$. For all $T = \prod_\alpha T_\alpha$ with $T_\alpha \subseteq \mathbf{Aut}(\mathcal{L}_\alpha)$, $\bigotimes_\alpha \mathcal{L}_\alpha$ and $\bigvee_\alpha \mathcal{L}_\alpha$ are in $\mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$. Moreover, $\bigotimes_\alpha \mathcal{L}_\alpha$ and $\bigvee_\alpha \mathcal{L}_\alpha$ are the bottom and the top elements of $\mathcal{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$ respectively, and $\mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$ and $\mathcal{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$, ordered by set-inclusion, are complete lattices.*

Proof. Let $\Sigma = \prod_\alpha \Sigma_\alpha$. Obviously, $\bigotimes_\alpha \mathcal{L}_\alpha$ and $\bigvee_\alpha \mathcal{L}_\alpha$ contain $\emptyset$ and Σ , and by definition $\bigotimes_\alpha \mathcal{L}_\alpha$ is $\cap$ -closed and $\bigvee_\alpha \mathcal{L}_\alpha$ contains all singletons. If $\omega \subseteq \bigvee_\alpha \mathcal{L}_\alpha$, then $(\cap \omega)_\beta[p] = \cap\{R_\beta[p]; R \in \omega\}$. Moreover, for all $p \in \Sigma$, $\cap_\alpha \pi_\alpha^{-1}(p_\alpha) = p$. As a consequence, $\bigotimes_\alpha \mathcal{L}_\alpha$ and $\bigvee_\alpha \mathcal{L}_\alpha$ are simple closure spaces on Σ , and obviously, Axioms P2 and P3 hold (note that for $A \subseteq \Sigma_\beta$, $p[A, \beta] \subseteq \cup_\alpha \pi_\alpha^{-1}(a_\alpha)$ if and only if $A \subseteq a_\beta$, or $p_\gamma \in a_\gamma$ for some $\gamma \neq \beta$, therefore $\vee_{\bigotimes} p[A, \beta] = p[\vee A, \beta]$).

Let $v \in T$, $a \in \prod_\alpha \mathcal{L}_\alpha$, $p \in \Sigma$, $R \subseteq \Sigma$, and $\omega \subseteq 2^\Sigma$. Then,

$$v(\cup_\alpha \pi_\alpha^{-1}(a_\alpha)) = \cup_\alpha \pi_\alpha^{-1}(v_\alpha(a_\alpha)),$$

and

$$v(R)_\beta[v(p)] = v_\beta(R_\beta[p]).$$

Moreover $v(\cap \omega) = \cap\{v(x); x \in \omega\}$. Hence, v is a bijection of $\bigotimes_\alpha \mathcal{L}_\alpha$ and of $\bigvee_\alpha \mathcal{L}_\alpha$, and v preserves arbitrary meets, hence also arbitrary joins.

Let $\mathcal{L} \in \mathcal{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$. By Axiom P2, $\bigotimes_\alpha \mathcal{L}_\alpha \subseteq \mathcal{L}$, whereas by Axiom P3, $\mathcal{L} \subseteq \bigvee_\alpha \mathcal{L}_\alpha$.

Finally, if $\omega \subseteq \mathcal{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$ (respectively $\omega \subseteq \mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$), then obviously $\cap \omega = \{a \in \mathcal{L}, \forall \mathcal{L} \in \omega\} \in \mathcal{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$ (respectively $\cap \omega \in \mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$). $\square$

Corollary 2.14. *Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$ and $\Sigma = \prod_\alpha \Sigma_\alpha$. Then*

$$\mathcal{S}(\mathcal{L}_\alpha, \alpha \in \Omega) = \{\mathcal{L} \in \mathbf{Cl}(\Sigma); \bigotimes_\alpha \mathcal{L}_\alpha \subseteq \mathcal{L} \subseteq \bigvee_\alpha \mathcal{L}_\alpha\}.$$

Remark 2.15. Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}\}_{\alpha \in \Omega}$. For all $T = \prod_\alpha T_\alpha$ with $T_\alpha \subseteq \mathbf{Aut}(\mathcal{L}_\alpha)$, $\mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$ is a complete meet-sublattice of $\mathcal{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$.

Proposition 2.16. *Let $\mathcal{L}_1, \mathcal{L}_2 \in \mathbf{Cl}$. Then $\mathcal{L}_1 \bigotimes \mathcal{L}_2 \cong \mathcal{L}_2 \bigotimes \mathcal{L}_1$, $\mathcal{L}_1 \bigvee \mathcal{L}_2 \cong \mathcal{L}_2 \bigvee \mathcal{L}_1$, and $2 \bigotimes \mathcal{L}_1 \cong \mathcal{L}_1 \cong 2 \bigvee \mathcal{L}_1$.*

Proof. Direct from Definition 2.11. $\square$

Proposition 2.17. *Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}\}_{\alpha \in \Omega}$ and $T = \prod_\alpha T_\alpha$ with $T_\alpha \subseteq \mathbf{Aut}(\mathcal{L}_\alpha)$. Then there is an isomorphism $f : \mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega) \rightarrow \mathcal{S}_T(2, \mathcal{L}_\alpha, \alpha \in \Omega)$ such that for all $\mathcal{L} \in \mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$, $\mathcal{L} \cong f(\mathcal{L})$.*

Proof. Direct from Definition 2.8. $\square$

Example 2.18. We now consider two examples, well known in many-body quantum physics, where instead of Axioms P1 and P2, we have

- (p1) $\prod_\alpha \Sigma_\alpha \subseteq \Sigma$,
- (p2) $(\prod_\alpha \Sigma_\alpha) \cap (\vee(\cup_\alpha \pi_\alpha^{-1}(a_\alpha))) = \cup_\alpha \pi_\alpha^{-1}(a_\alpha), \forall a \in \prod_\alpha \mathcal{L}_\alpha$,

Let $\mathcal{H}_1$ and $\mathcal{H}_2$ be complex Hilbert spaces, $\mathcal{L}_1 = \mathbf{P}(\mathcal{H}_1)$, $\mathcal{L}_2 = \mathbf{P}(\mathcal{H}_2)$ and $\mathcal{L} = \mathbf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$. Then, Axioms p1, p2 and P3 hold in $\mathcal{L}$. Moreover, Axiom P4 holds for $T = \mathbf{U}(\mathcal{H}_1) \times \mathbf{U}(\mathcal{H}_2)$. Note that $\vee(A_1 \times A_2) = 1$, for all $A_i \subseteq \Sigma_{\mathcal{H}_i}$ with $\vee A_i = 1$.

Let $\mathcal{H}$ be a complex Hilbert space and $\mathcal{F} = \bigoplus_{n \geq 0} \mathcal{H}^{\otimes n}$ be the Fock space (neither symmetrized, nor antisymmetrized). Let $\mathcal{L} = \mathbf{P}(\mathcal{F})$, and for all integer i , let $\mathcal{L}_i = \mathbf{P}(\mathcal{H})$. Let n be an integer. Consider the family $\{\mathcal{L}_i; 1 \leq i \leq n\}$. Then Axioms p1, p2, P3 and P4 with $T_i = \mathbf{U}(\mathcal{H})$ hold in $\mathcal{L}$ for all n . For all i , let $\Sigma_i = \Sigma_{\mathcal{H}}$. Note that in that case, for all n we have $\vee(\prod_{i=1}^n \Sigma_i) \neq 1$.

3. COMPARISON WITH OTHER TENSOR PRODUCTS

In this section, we compare the bottom element $\mathcal{L}_1 \bigcirc \mathcal{L}_2$ with the *separated product* of Aerts, with the *box product* $\mathcal{L}_1 \square \mathcal{L}_2$ of Grätzer and Wehrung, and with the *lattice tensor product* $\mathcal{L}_1 \boxtimes \mathcal{L}_2$. On the other hand, we compare the top element $\mathcal{L}_1 \bigvee \mathcal{L}_2$ with the semilattice tensor product of Fraser and with the tensor products of Chu, Golfin and Shmueli.

3.1. The separated product.

Definition 3.1 (D. Aerts, [1]). Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$ with for all $\alpha \in \Omega$, $\mathcal{L}_\alpha$ orthocomplemented, $\Sigma = \prod_{\alpha} \Sigma_\alpha$, and let $p, q \in \Sigma$. Denote by $\#$ the binary relation on Σ defined by $p \# q$ if and only if there is $\beta \in \Omega$ such that $p_\beta \perp_\beta q_\beta$. Then

$$\bigotimes_{\text{Aerts}} \mathcal{L}_\alpha := \{R \subseteq \Sigma; R^{\# \#} = R\}.$$

Lemma 3.2. Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$ with for all $\alpha \in \Omega$, $\mathcal{L}_\alpha$ orthocomplemented. Then

$$\bigotimes_{\text{Aerts}} \mathcal{L}_\alpha = \bigcirc_{\alpha} \mathcal{L}_\alpha,$$

and $\bigotimes_{\text{Aerts}} \mathcal{L}_\alpha$ is orthocomplemented.

Proof. Let $\Sigma = \prod_{\alpha} \Sigma_\alpha$. Obviously, $\#$ is symmetric and anti-reflexive. Since $\mathcal{L}_\alpha$ is orthocomplemented, $\perp_\alpha$ is *separating*, i.e. for all $p_\alpha, q_\alpha \in \Sigma_\alpha$, there is $r_\alpha \in \Sigma_\alpha$ such that $p_\alpha \perp_\alpha r_\alpha$ and $q_\alpha \not\perp_\alpha r_\alpha$. Therefore, it follows directly from Definition 3.1 that $\#$ is separating. As a consequence, $\bigotimes_{\text{Aerts}} \mathcal{L}_\alpha$ is an orthocomplemented simple closure space on Σ . Moreover, coatoms are given by $p^\# = \cup_{\alpha} \pi_{\alpha}^{-1}(p_{\alpha}^{\perp_{\alpha}})$. As a consequence, $\bigotimes_{\text{Aerts}} \mathcal{L}_\alpha \subseteq \bigcirc_{\alpha} \mathcal{L}_\alpha$.

Let $a \in \prod_{\alpha} \mathcal{L}_\alpha$. Denote the set of coatoms above a_α by $\Sigma'[a_\alpha]$. Then $a_\alpha = \cap_{\alpha} \Sigma'[a_\alpha]$. Moreover,

$$\bigcap \{\cup_{\alpha} \pi_{\alpha}^{-1}(x_{\alpha}); x \in \prod_{\alpha} \Sigma'[a_{\alpha}]\} = \cup_{\alpha} \pi_{\alpha}^{-1}(a_{\alpha}).$$

As a consequence, $\bigcirc_{\alpha} \mathcal{L}_\alpha \subseteq \bigotimes_{\text{Aerts}} \mathcal{L}_\alpha$. □

Remark 3.3. The symbol $\bigcirc$ was originally used by Aerts.

3.2. The box product and the lattice tensor product.

Definition 3.4 (G. Grätzer, F. Wehrung, [9]). Let $\mathcal{L}_1$ and $\mathcal{L}_2$ be lattices and $(a, b) \in \mathcal{L}_1 \times \mathcal{L}_2$. Define $a \square b = \{(x, y) \in \mathcal{L}_1 \times \mathcal{L}_2; x \leq a \text{ or } y \leq b\}$. The *box product* $\mathcal{L}_1 \square \mathcal{L}_2$ is defined as the set of all finite intersections of the form $\cap \{a_i \square b_i; i \leq n\}$ where $(a_i, b_i) \in \mathcal{L}_1 \times \mathcal{L}_2$ for all $i \leq n$, ordered by set-inclusion ([9], Definition 2.1).

Define $a \boxtimes b = \{(x, y) \in \mathcal{L}_1 \times \mathcal{L}_2; x \leq a \text{ and } y \leq b\} \cup (\mathcal{L}_1 \times \perp_{\mathcal{L}_2}) \cup (\perp_{\mathcal{L}_1} \times \mathcal{L}_2)$ where $\perp_{\mathcal{L}_i} = \{0_i\}$ if $\mathcal{L}_i$ has a zero, and $\emptyset$ otherwise. Call a subset X of $\mathcal{L}_1 \times \mathcal{L}_2$

confined if $X \subseteq a \boxtimes b$ for some $(a, b) \in \mathcal{L}_1 \times \mathcal{L}_2$. Let $\mathcal{L}_1 \boxtimes \mathcal{L}_2$ be the set of all confined elements of $\mathcal{L}_1 \square \mathcal{L}_2$, ordered by set-inclusion. If $\mathcal{L}_1 \boxtimes \mathcal{L}_2$ is nonempty, then it is called the *the lattice tensor product* of $\mathcal{L}_1$ and $\mathcal{L}_2$ (see [9], Section 3).

Definition 3.5. Let $\mathcal{L}$ be a lattice and $a \in \mathcal{L}$. We denote by $a \downarrow$ the set $\{x \in \mathcal{L}; x \leq a\}$ and by L the set $\{a \downarrow; a \in \mathcal{L}\}$ ordered by set-inclusion (note that $L \subseteq 2^{\mathcal{L}}$).

Let Σ_1 and Σ_2 be nonempty sets, $\mathcal{L}_1 \subseteq 2^{\Sigma_1}$ and $\mathcal{L}_2 \subseteq 2^{\Sigma_2}$. We define $\mathcal{L}_1 \otimes_n \mathcal{L}_2$ by taking only finite intersections in Definition 2.11. If $\mathcal{L}_1$ and $\mathcal{L}_2$ are atomistic lattices, then $\mathcal{L}_1 \otimes_n \mathcal{L}_2$ is defined as $l_1 \otimes_n l_2$, where $l_i = \{\Sigma[a_i]; a_i \in \mathcal{L}_i\} \subseteq 2^{\Sigma_i}$ and $\Sigma[a_i]$ denotes the set of atoms under a_i ($\Sigma_i := \Sigma[1_i]$).

Proposition 3.6. Let $\mathcal{L}_1$ and $\mathcal{L}_2$ be lattices with 1. Then $\mathcal{L}_1 \square \mathcal{L}_2 = L_1 \otimes_n L_2$.

Proof. Direct from Definitions 3.4 and 3.5. $\square$

Lemma 3.7 ([9], Corollary 4.2). Let $\mathcal{L}_1$ and $\mathcal{L}_2$ be lattices with 0. Then $\mathcal{L}_1^* \square \mathcal{L}_2^* \cong (\mathcal{L}_1 \boxtimes \mathcal{L}_2)^*$.

Theorem 3.8. For atomistic lattices with 1 (and 0), $\mathcal{L}_1 \otimes_n \mathcal{L}_2 \cong \mathcal{L}_1 \square \mathcal{L}_2$, and in case $\mathcal{L}_1$ and $\mathcal{L}_2$ are moreover coatomistic, $\mathcal{L}_1 \otimes_n \mathcal{L}_2 \cong \mathcal{L}_1 \boxtimes \mathcal{L}_2$.

Proof. Define $f : \mathcal{L}_1 \otimes_n \mathcal{L}_2 \rightarrow \mathcal{L}_1 \square \mathcal{L}_2$ as

$$f(\Sigma[a_1] \times \Sigma_2 \cup \Sigma_1 \times \Sigma[a_2]) = a_1 \square a_2 := (a_1 \downarrow) \times \mathcal{L}_2 \cup \mathcal{L}_1 \times (a_2 \downarrow),$$

and $f(\cap x_i) := \cap f(x_i)$. Obviously, f is bijective and preserves meets, hence also joins.

From Lemma 3.7 and what precedes, we have $(\mathcal{L}_1 \boxtimes \mathcal{L}_2)^* \cong \mathcal{L}_1^* \otimes_n \mathcal{L}_2^*$. Now, an atom of $\mathcal{L}_1^* \otimes_n \mathcal{L}_2^*$ is of the form $r = \{(x_1, x_2)\}$ with x_1 and x_2 coatoms, and a coatom is of the form $x = \Sigma'[p_1] \times \Sigma'_2 \cup \Sigma'_1 \times \Sigma'[p_2]$ with p_1 and p_2 atoms and where $\Sigma'[a]$ denotes the set of coatoms above a ($\Sigma'_i := \Sigma'[0_i]$). Moreover, $r \leq x$ if and only if $p_1 \leq x_1$ or $p_2 \leq x_2$.

On the other hand, an atom of $(\mathcal{L}_1 \otimes_n \mathcal{L}_2)^*$ is of the form $s = \Sigma[x_1] \times \Sigma_2 \cup \Sigma_1 \cup \Sigma[x_2]$ and a coatom is of the form $y = \{(p_1, p_2)\}$, and $s \leq y$ if and only if $p_1 \leq x_1$ or $p_2 \leq x_2$. Therefore, there is obviously an isomorphism between $\mathcal{L}_1^* \otimes_n \mathcal{L}_2^*$ and $(\mathcal{L}_1 \otimes_n \mathcal{L}_2)^*$, hence between $\mathcal{L}_1 \boxtimes \mathcal{L}_2$ and $\mathcal{L}_1 \otimes_n \mathcal{L}_2$. $\square$

3.3. The tensor products of Chu, Golfin and Shmuely.

Definition 3.9 (P. H. Chu, [3] §II.1, III.3). The category **Chu**(Set, 2) has as objects triples $\mathbf{A} = (A, r, X)$ where A and X are sets and r is a map from $A \times X$ to 2 (where $2 = \{0, 1\}$), and as arrows pairs of maps $(F, G) : \mathbf{A} \rightarrow (B, s, Y)$ with $F : A \rightarrow B$ and $G : Y \rightarrow X$ such that $s(F(a), y) = r(a, G(y))$ for all $a \in A$ and $y \in Y$. There is a functor $^\perp$ and a bifunctor $\otimes_{\text{Chu}}$ defined on objects as $\mathbf{A}^\perp = (X, \check{r}, A)$ with $\check{r}(x, a) = r(a, x)$, and $\mathbf{A} \otimes_{\text{Chu}} \mathbf{B} = (A \times B, t, \mathbf{Chu}(\mathbf{A}, \mathbf{B}^\perp))$ with $t((a, b), (F, G)) = r(a, G(b)) = \check{s}(F(a), b)$.

Remark 3.10. A simple closure space $\mathcal{L} \in \mathbf{Cl}(\Sigma)$ is of course a Chu space, which we also denote by $\mathcal{L}$ (for $p \in \Sigma$ and $a \in \mathcal{L}$, put $r(p, a) = 1$ if and only if $p \in a$).

Proposition 3.11. Let $\mathcal{L}_0 \in \mathbf{Cl}(\Sigma_0)$ and $\mathcal{L}_1 \in \mathbf{Cl}(\Sigma_1)$. Then there is a one to one correspondence between invertible arrows in **Chu**($\mathcal{L}_0, \mathcal{L}_1$) and isomorphisms between $\mathcal{L}_0$ and $\mathcal{L}_1$.

Proof. Let $f : \mathcal{L}_0 \rightarrow \mathcal{L}_1$ be an isomorphism. Then $(f, f^{-1}) \in \mathbf{Chu}(\mathcal{L}_0, \mathcal{L}_1)$. Let (F, G) be an invertible arrow in $\mathbf{Chu}(\mathcal{L}_0, \mathcal{L}_1)$. Then $G : \mathcal{L}_1 \rightarrow \mathcal{L}_0$ is bijective, and since $G = F^{-1}$, G preserves arbitrary meets, hence also arbitrary joins. $\square$

Definition 3.12. Let $\mathcal{L}_1$ and $\mathcal{L}_2$ be posets. A Galois connection between $\mathcal{L}_1$ and $\mathcal{L}_2$ is a pair (f, g) of order-preserving maps with $f : \mathcal{L}_1 \rightarrow \mathcal{L}_2$ and $g : \mathcal{L}_2 \rightarrow \mathcal{L}_1$, such that for all $a \in \mathcal{L}_1$ and $b \in \mathcal{L}_2$, $f(a) \leq b \Leftrightarrow a \leq g(b)$.

Lemma 3.13. Let $\mathcal{L}_0$ and $\mathcal{L}_1$ be complete join-semilattices and (f, g) a pair of maps with $f : \mathcal{L}_0 \rightarrow \mathcal{L}_1$ and $g : \mathcal{L}_1 \rightarrow \mathcal{L}_0$.

- (1) If (f, g) forms a Galois connection, then f preserves arbitrary joins.
- (2) Suppose that $\mathcal{L}_0 \in \mathbf{Cl}(\Sigma_0)$. For $b \in \mathcal{L}_1$, let $F^{-1}(b)$ denote the set $\{p \in \Sigma_0 ; f(p) \leq b\}$. Then f preserves arbitrary joins $\Leftrightarrow f(a) = \vee\{f(p) ; p \in a\}$ for all $a \in \mathcal{L}_0$ and $F^{-1}(b) \in \mathcal{L}_0$ for all $b \in \mathcal{L}_1$.

Proof. (1) Let $\omega \subseteq \mathcal{L}_0$. Since f preserves order, $y := \vee\{f(x) ; x \in \omega\} \leq f(\vee\omega)$. On the other hand, for any $x \in \omega$, $f(x) \leq y$, hence $x \leq g(y)$. As a consequence, $\vee\omega \leq g(y)$, therefore $f(\vee\omega) \leq y$.

(2) $(\Rightarrow)$ If $p \in \vee F^{-1}(b)$, then $f(p) \leq \vee\{f(q) ; q \in F^{-1}(b)\} \leq b$, hence $p \in F^{-1}(b)$. $(\Leftarrow)$ Obviously, (f, F^{-1}) forms a Galois connection between $\mathcal{L}_0$ and $\mathcal{L}_1$. $\square$

Theorem 3.14. Let $\mathcal{L}_1 \in \mathbf{Cl}(\Sigma_1)$ and $\mathcal{L}_2 \in \mathbf{Cl}(\Sigma_2)$. Then,

- (1) $\mathcal{L}_1 \odot \mathcal{L}_2 \cong \mathcal{L}_1 \otimes_{\mathbf{Chu}} \mathcal{L}_2$,
- (2) $(\mathcal{L}_1 \odot \mathcal{L}_2)^* \cong \{f : \mathcal{L}_1 \rightarrow \mathcal{L}_2^* ; f \text{ preserves arbitrary joins}\}$ ordered pointwise (i.e. $f \leq g$ if and only if $f(p_1) \leq_* g(p_1)$ for all $p_1 \in \Sigma_1$, where $\leq_*$ denotes the order relation in $\mathcal{L}_2^*$).

Proof. (1) We denote $\Sigma_1 \times \Sigma_2$ by Σ . Let $(F, G) \in \mathbf{Chu}(\mathcal{L}_1, \mathcal{L}_2^\perp)$. Define

$$R_{(F,G)} := \bigcup_{p_1 \in \Sigma_1} p_1 \times F(p_1).$$

By definition, $R_{(F,G)_2}[p] \in \mathcal{L}_2$ for all $p \in \Sigma$, and

$$q_1 \in R_{(F,G)_1}[p] \Leftrightarrow (q_1, p_2) \in R_{(F,G)} \Leftrightarrow p_2 \in F(q_1) \Leftrightarrow q_1 \in G(p_2),$$

thus $R_{(F,G)_1}[p] \in \mathcal{L}_1$ for all $p \in \Sigma$, hence $R_{(F,G)} \in \mathcal{L}_1 \odot \mathcal{L}_2$.

Let $R \in \mathcal{L}_1 \odot \mathcal{L}_2$. Define $F_R : \Sigma_1 \rightarrow \mathcal{L}_2$ and $G_R : \Sigma_2 \rightarrow \mathcal{L}_1$ as $F_R(p_1) := R_2[(p_1, \cdot)]$ and $G_R(p_2) := R_1[(\cdot, p_2)]$. Then

$$p_2 \in F_R(p_1) \Leftrightarrow (p_1, p_2) \in R \Leftrightarrow p_1 \in G_R(p_2),$$

so that $(F_R, G_R) \in \mathbf{Chu}(\mathcal{L}_1, \mathcal{L}_2^\perp)$.

Obviously,

$$F_{R_{(F,G)}} = F, \quad G_{R_{(F,G)}} = G, \quad \text{and} \quad R_{(F_R, G_R)} = R.$$

As a consequence, there is an invertible arrow in $\mathbf{Chu}(\mathcal{L}_1 \odot \mathcal{L}_2, \mathcal{L}_1 \otimes_{\mathbf{Chu}} \mathcal{L}_2)$.

(2) Let $(F, G) \in \mathbf{Chu}(\mathcal{L}_1, \mathcal{L}_2^\perp)$. Define $f : \mathcal{L}_1 \rightarrow \mathcal{L}_2^*$ and $g : \mathcal{L}_2^* \rightarrow \mathcal{L}_1$ as $f(a) := \cap F(a)$ and $g(b) = \cap G(b)$. Then, obviously (f, g) forms a Galois connection between $\mathcal{L}_1$ and $\mathcal{L}_2^*$.

Let $f : \mathcal{L}_1 \rightarrow \mathcal{L}_2^*$ be $\vee$ -preserving. Define $g : \mathcal{L}_2 \rightarrow \mathcal{L}_1$ as $g(b) := \vee\{a \in \mathcal{L}_1 ; f(a) \leq_* b\}$. Then, obviously $(f, g) \in \mathbf{Chu}(\mathcal{L}_1, \mathcal{L}_2^\perp)$. $\square$

Remark 3.15. Note that $\otimes$ is the $\boxtimes$ -tensor product of Golfin ([8], Definition II.3.6, II.1.1). As a corollary of Theorem 3.14 part 2, $\mathcal{L}_1 \otimes \mathcal{L}_2 \cong \mathcal{L}_1 \otimes \mathcal{L}_2$ the tensor product of Shmueli [19]. Note that there is a one to one correspondence between maps from $\mathcal{L}_1$ to $\mathcal{L}_2^*$ preserving arbitrary joins and Galois connections between $\mathcal{L}_1$ and $\mathcal{L}_2^*$.

3.4. The semilattice tensor product of Fraser.

Definition 3.16. Let $\mathbf{C}$ be a concrete category over $\mathbf{Set}$. Let $\{A_\alpha\}_{\alpha \in \Omega}$ and C be objects of $\mathbf{C}$. A *multi-morphism* of $\mathbf{C}$ is an arrow $f \in \mathbf{Set}(\prod_{\alpha} A_\alpha, C)$ such that for all $a \in \prod_{\alpha} A_\alpha$, $f(a[-, \alpha]) \in C(A_\alpha, C)$, where for $x \in A_\alpha$, $a[x, \alpha]$ is defined as $\pi_\alpha(a[x, \alpha]) = x$ and $\pi_\beta(a[x, \alpha]) = a_\beta$, for all $\beta \neq \alpha$. If $|\Omega| = 2$, then f is called a *bimorphism*.

Definition 3.17 (G. Seal, [18]). Let $\mathbf{C}$ be a concrete category over $\mathbf{Set}$. A *tensor product* in $\mathbf{C}$ is a bifunctor $- \otimes - : \mathbf{C} \times \mathbf{C} \rightarrow \mathbf{C}$ such that for all objects A, B of $\mathbf{C}$, there is a bimorphism $f : A \times B \rightarrow A \otimes B$, and for all object C of $\mathbf{C}$, and all bimorphism $g : A \times B \rightarrow C$, there is a unique arrow $h \in C(A \otimes B, C)$ which makes the diagram

$$\begin{array}{ccc} A \times B & \xrightarrow{f} & A \otimes B \\ \downarrow g & \searrow !h & \\ C & & \end{array}$$

commute.

Remark 3.18. By definition, the tensor product is unique up to isomorphisms. For the category of join-semilattices with maps preserving finite joins, Definition 3.17 is equivalent to the definition of join-semilattice tensor product given by Fraser in [7] (note that $f(\mathcal{L}_1 \times \mathcal{L}_2)$ generates $\mathcal{L}_1 \otimes \mathcal{L}_2$ if and only if the arrow h is unique). Therefore, we call the tensor product in the category of complete join-semilattices with maps preserving arbitrary joins, the complete join-semilattice tensor product. For $\mathcal{L}_1, \mathcal{L}_2 \in \mathbf{Cl}$, it is given by $\mathcal{L}_1 \otimes \mathcal{L}_2$ (Theorem 3.20).

Lemma 3.19. Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$, and $\mathcal{L} \in \mathbf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$. Then, for all $a \in \prod_{\alpha} \mathcal{L}_\alpha$, $\prod_{\alpha} a_\alpha \in \mathcal{L}$. Moreover, the map $f : \prod_{\alpha} \mathcal{L}_\alpha \rightarrow \mathcal{L}$ defined as $f(a) = \prod_{\alpha} a_\alpha$, is an injective multi-morphism of $\mathbf{Cl}$.

Proof. Obviously $\prod_{\alpha} a_\alpha = \cap_{\alpha} \pi^{-1}(a_\alpha)$, hence the first statement follows from Axiom P2.

Let $\Sigma = \prod_{\alpha} \Sigma_\alpha$, $p \in \Sigma$, $\beta \in \Omega$, $\omega \subseteq \mathcal{L}_\beta$, and $a \in \prod_{\alpha} \mathcal{L}_\alpha$. By Axiom P3,

$$\bigvee_{x \in \omega} p[x, \beta] = p[\bigvee \omega, \beta].$$

As a consequence,

$$\bigvee_{x \in \Omega} f(a[x, \beta]) = \bigvee_{x \in \omega} \bigvee_{p \in \prod_{\alpha} a_\alpha} p[x, \beta] = \bigvee_{p \in \prod_{\alpha} a_\alpha} p[\bigvee \omega, \beta] = f(a[\bigvee \omega, \beta]).$$

□

Theorem 3.20. $\otimes$ is a tensor product in $\mathbf{Cl}$.

Proof. Let $\mathcal{L}_1, \mathcal{L}_2 \in \mathbf{Cl}$. By Theorem 2.13, $\mathcal{L}_1 \odot \mathcal{L}_2 \in \mathbf{Cl}$. By Lemma 3.19, the map $f : \mathcal{L}_1 \times \mathcal{L}_2 \rightarrow \mathcal{L}_1 \odot \mathcal{L}_2$ defined as $f(a_1, a_2) := a_1 \times a_2$ is a bimorphism.

Let $\mathcal{L} \in \mathbf{Cl}$ (or a complete join-semilattice), and let $g : \mathcal{L}_1 \times \mathcal{L}_2 \rightarrow \mathcal{L}$ be a bimorphism. Define $h : \mathcal{L}_1 \odot \mathcal{L}_2 \rightarrow \mathcal{L}$ by $h(R) = \vee \{g(p_1, p_2) ; (p_1, p_2) \in R\}$. Let $b \in \mathcal{L}$. Then (see Lemma 3.13 for notations),

$$H^{-1}(b) = \bigcup_{p_1 \in \Sigma_1} p_1 \times G_{p_1}^{-1}(b) = \bigcup_{p_2 \in \Sigma_2} G_{p_2}^{-1}(b) \times p_2,$$

where $g_{p_1} := g(p_1, -)$ and $g_{p_2} := g(-, p_2)$. Since g is a bimorphism, it follows from Lemma 3.13 and Definition 2.11 that $H^{-1}(b) \in \mathcal{L}_1 \odot \mathcal{L}_2$. Therefore, by Lemma 3.13, $h \in \mathbf{Cl}(\mathcal{L}_1 \odot \mathcal{L}_2, \mathcal{L})$. Finally, if $h' \in \mathbf{Cl}(\mathcal{L}_1 \odot \mathcal{L}_2, \mathcal{L})$ and $h' \circ f = g$, then h' equals h on atoms, therefore $h' = h$.

Let $L_1, L_2 \in \mathbf{Cl}$, $f_1 \in \mathbf{Cl}(\mathcal{L}_1, L_1)$, and $f_2 \in \mathbf{Cl}(\mathcal{L}_2, L_2)$. Then $g = f \circ (f_1 \times f_2) : \mathcal{L}_1 \times \mathcal{L}_2 \rightarrow L_1 \odot L_2$ is a bimorphism. We define $f_1 \odot f_2 \in \mathbf{Cl}(\mathcal{L}_1 \odot \mathcal{L}_2, L_1 \odot L_2)$ to be the arrow h constructed above. $\square$

4. EQUIVALENT DEFINITION

In this section we give an equivalent definition of $\mathbf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$ and $\mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$ in terms of a universal property with respect to a given class of bimorphisms of $\mathbf{Cl}$.

Lemma 4.1. *Let $\mathcal{L} \in \mathbf{Cl}(\Sigma)$. For all $a \in \mathcal{L}$, there is (a unique) $h_a : \mathcal{L} \rightarrow 2$ preserving arbitrary joins such that for all $p \in \Sigma$, $h_a(p) = 0$ if and only if $p \in a$.*

Proof. Let $b \in \mathcal{L}$. Define h_a by $h_a(b) = \vee \{h_a(p) ; p \in b\}$ (hence $h_a(0) = 0$) with $h_a(p) = 0$ if $p \in a$ and $h_a(p) = 1$ if $p \notin a$. Then, by Lemma 3.13, h preserves arbitrary joins. $\square$

Lemma 4.2. *Let $\mathcal{L} \in \mathbf{Cl}$, $\{\mathcal{L}_\alpha \in \mathbf{Cl}\}_{\alpha \in \Omega}$, $T = \prod_{\alpha} T_\alpha$ with $T_\alpha \subseteq \mathbf{Aut}(\mathcal{L}_\alpha)$, and $f : \prod_{\alpha} \mathcal{L}_\alpha \rightarrow \mathcal{L}$ a multi-morphism of $\mathbf{Cl}$. Define*

$$V_T(f) := \{f \circ v : \prod_{\alpha} \mathcal{L}_\alpha \rightarrow \mathcal{L} ; v \in T\},$$

$$\Gamma_{\cap} := \{\cap_{\alpha} g_{\alpha} : \prod_{\alpha} \mathcal{L}_\alpha \rightarrow 2 ; g_{\alpha} \in \mathbf{Cl}(\mathcal{L}_\alpha, 2)\},$$

with $\cap_{\alpha} g_{\alpha}(a) := \cap \{g_{\alpha}(a_{\alpha}) ; \alpha \in \Omega\}$. Then the maps in $V_T(f)$ and $\Gamma_{\cap}$ are multi-morphisms of $\mathbf{Cl}$.

Proof. Let $g = \cap_{\alpha} g_{\alpha} \in \Gamma_{\cap}$, $a \in \prod_{\alpha} \mathcal{L}_\alpha$, $\beta \in \Omega$, and $b \in \mathcal{L}$. Write $h(-) := g(a[-, \beta]) : \mathcal{L}_{\beta} \rightarrow 2$. Then,

$$H^{-1}(0) := \{p_{\beta} \in \Sigma_{\beta} ; g_{\beta}(p_{\beta}) \bigcap_{\alpha \neq \beta} g_{\alpha}(a_{\alpha}) = 0\}$$

equals 1 if $g_{\alpha}(a_{\alpha}) = 0$ for some $\alpha \neq \beta$, and $G_{\beta}^{-1}(0)$ otherwise. Therefore, by Lemma 3.13, h preserves arbitrary joins, hence g is a multi-morphism of $\mathbf{Cl}$.

On the other hand, let $w = f \circ v \in V_T$ and $\omega \subseteq \mathcal{L}_{\beta}$. Then

$$w(a[\vee \omega, \beta]) = f(v(a)[\vee v_{\beta}(x) ; x \in \omega]) = \bigvee_{x \in \omega} w(a[x, \beta]),$$

since f is a multi-morphism. $\square$

Definition 4.3. Let $\mathcal{L} \in \mathbf{Cl}$ and $A \subseteq \mathcal{L}$. We say that A *generates* $\mathcal{L}$ if each element of $\mathcal{L}$ is the join of elements in A .

Theorem 4.4. *Let $\mathcal{L} \in \mathbf{Cl}(\Sigma)$, $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$, and $T = \prod_\alpha T_\alpha$ with $T_\alpha \subseteq \text{Aut}(\mathcal{L}_\alpha)$.*

- (1) *If $\mathcal{L} \in \mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$, then there is a multi-morphism $f : \prod_\alpha \mathcal{L}_\alpha \rightarrow \mathcal{L}$ of $\mathbf{Cl}$ such that $f(\prod_\alpha \mathcal{L}_\alpha)$ generates $\mathcal{L}$, and for all $g \in \Gamma_\cap$ and $w \in V_T(f)$, there is a unique $h \in \mathbf{Cl}(\mathcal{L}, 2)$ and a unique $u \in \mathbf{Cl}(\mathcal{L}, \mathcal{L})$ such that the diagrams*

$$\begin{array}{ccc} \prod_\alpha \mathcal{L}_\alpha & \xrightarrow{f} & \mathcal{L} \\ g \downarrow & \nearrow !h & \\ 2 & & \end{array} \quad \begin{array}{ccc} \prod_\alpha \mathcal{L}_\alpha & \xrightarrow{f} & \mathcal{L} \\ w \downarrow & \nearrow !u & \\ \mathcal{L} & & \end{array}$$

commute.

- (2) *Conversely, if there is a multi-morphism of $\mathbf{Cl}$ $f : \prod_\alpha \mathcal{L}_\alpha \rightarrow \mathcal{L}$ satisfying all conditions of part (1), then there is $\mathcal{L}_0 \in \mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$ such that $\mathcal{L} \cong \mathcal{L}_0$.*

Proof. (1) By Lemma 3.19, the map $f : \prod_\alpha \mathcal{L}_\alpha \rightarrow \mathcal{L}$ defined as $f(a) = \prod_\alpha a_\alpha$, is a multi-morphism of $\mathbf{Cl}$, and obviously, $f(\prod_\alpha \mathcal{L}_\alpha)$ generates $\mathcal{L}$. Moreover, by Axiom P4, for all $w \in V_T(f)$ there is a unique u such that the second diagram commutes.

Let $g = \cap_\alpha g_\alpha \in \Gamma_\cap$ and $\Sigma = \prod_\alpha \Sigma_\alpha$. Define $a_\alpha := \vee G_\alpha^{-1}(0)$ and $x := \cup_\alpha \pi_\alpha^{-1}(a_\alpha)$ (see Lemma 3.13). By Axiom P2 $x \in \mathcal{L}$, and by Lemma 4.1 there is $h_x \in \mathbf{Cl}(\mathcal{L}, 2)$ such that for all $p \in \Sigma$, $h_x(p) = 0$ if and only if $p \in x$. Hence, $h_x \circ f = g$. Let $h \in \mathbf{Cl}(\mathcal{L}, 2)$ such that $h \circ f = g$. Then on atoms h equals h_x , therefore $h = h_x$.

(2) Let $\Sigma_0 = \prod_\alpha \Sigma_\alpha$. We denote by F the mapping from Σ_0 to $\mathcal{L}$ induced by the multi-morphism f .

(2.1) For $a, b \in \prod_\alpha \mathcal{L}_\alpha$, we write $a \leq b$ if and only if $a_\beta \leq b_\beta$, for all $\beta \in \Omega$.

Claim: The following implication holds: $f(a) \subseteq f(b) \Rightarrow a \leq b$. As a corollary, f is injective. [*Proof:* Suppose that $f(a) \subseteq f(b)$ and that $a_\beta \not\leq b_\beta$ for some $\beta \in \Omega$. By Lemma 4.1, there is $h_{b_\beta} : \mathcal{L}_\beta \rightarrow 2$ such that for all $p \in \Sigma_\beta$, $h_{b_\beta}(p) = 0$ if and only if $p \in b_\beta$. Let $g_\beta := h_{b_\beta}$, $g_\alpha := h_{0_\alpha}$ for all $\alpha \neq \beta$, and $g := \cap_\alpha g_\alpha$. By definition, $g \in \Gamma_\cap$, hence there is $h \in \mathbf{Cl}(\mathcal{L}, 2)$ such that $h \circ f = g$. As a consequence,

$$1 = g_\beta(a_\beta) = g(a) = h(f(a)) \subseteq h(f(b)) = g(b) = g_\beta(b_\beta) = 0,$$

a contradiction.]

As a consequence, since $f(\prod_\alpha \mathcal{L}_\alpha)$ generates $\mathcal{L}$, for all $p \in \Sigma_0$, $F(p)$ is an atom of $\mathcal{L}$, and the mapping from Σ_0 to Σ induced by F (which we also denote by F) is bijective. Moreover, for all $a \in \prod_\alpha \mathcal{L}_\alpha$,

$$f(a) = \vee \{F(p) ; p \in \prod_\alpha a_\alpha\}.$$

Therefore, for all $a, b \in \prod_\alpha \mathcal{L}_\alpha$, we have: $a \leq b \Rightarrow f(a) \subseteq f(b)$ (note that if Ω is finite, then this implication follows directly from the fact that f is a multi-morphism).

(2.2) Let $\mathcal{L}_0 \subseteq 2^{\Sigma_0}$ defined as $\mathcal{L}_0 := \{F^{-1}(c) ; c \in \mathcal{L}\}$. Then by what precedes, $\mathcal{L}_0 \in \mathbf{Cl}(\Sigma_0)$ and the map $F^{-1} : \mathcal{L} \rightarrow \mathcal{L}_0$ is bijective and preserves arbitrary meets, hence also arbitrary joins. It remains to prove that $\mathcal{L}_0 \in \mathcal{S}_T(\mathcal{L}_\alpha, \alpha \in \Omega)$, hence to check that Axioms P2, P3 and P4 hold in $\mathcal{L}_0$. Below, if $g \in \Gamma_\cap$, then G denotes the map from Σ_0 to 2 induced by g .

(P2) Let $a \in \prod_{\alpha} \mathcal{L}_{\alpha}$, $x := F(\cup_{\alpha} \pi_{\alpha}^{-1}(a_{\alpha}))$, and $p \in \Sigma_0$ such that $p_{\alpha} \notin a_{\alpha}$ for all $\alpha \in \Omega$. Suppose that $F(p) \in \vee x$. From Lemma 4.1, for all $\beta \in \Omega$, there is $h_{a_{\beta}} \in \mathbf{Cl}(\mathcal{L}_{\beta}, 2)$ such that for all $p \in \Sigma_{\beta}$, $h_{a_{\beta}}(p) = 0$ if and only if $p \in a_{\beta}$. Let $g = \cap_{\alpha} h_{a_{\alpha}}$. By definition, $g \in \Gamma_{\cap}$, hence there is $h \in \mathbf{Cl}(\mathcal{L}, 2)$ such that $h \circ f = g$; whence

$$\begin{aligned} 1 = G(p) &= h(F(p)) \subseteq h(\vee x) = \vee \{h(F(p)) ; p \in \cup_{\alpha} \pi_{\alpha}^{-1}(a_{\alpha})\} \\ &= \vee \{G(p) ; p \in \cup_{\alpha} \pi_{\alpha}^{-1}(a_{\alpha})\} = 0, \end{aligned}$$

a contradiction. As a consequence, $F^{-1}(\vee F(\cup_{\alpha} \pi_{\alpha}^{-1}(a_{\alpha}))) = \cup_{\alpha} \pi_{\alpha}^{-1}(a_{\alpha})$, hence $\cup_{\alpha} \pi_{\alpha}^{-1}(a_{\alpha}) \in \mathcal{L}_0$.

(P3) Let $\beta \in \Omega$, $A \subseteq \Sigma_{\beta}$, and $p \in \Sigma_0$, such that $p[A, \beta] \in \mathcal{L}_0$, i.e. there is $c \in \mathcal{L}$ such that $p[A, \beta] = F^{-1}(c)$. Let $q \in \vee A$ and let $\hat{p} \in \prod_{\alpha} \mathcal{L}_{\alpha}$ such that $\prod_{\alpha} \hat{p}_{\alpha} = \{p\}$. Then $p[q, \beta] \in p[\vee A, \beta]$, hence, since f is multi-morphism, we find that

$$F(p[q, \beta]) \in F(p[\vee A, \beta]) \subseteq \vee F(p[\vee A, \beta]) = f(\hat{p}[\vee A, \beta]) = \bigvee_{q \in A} F(p[q, \beta]) \subseteq c.$$

As a consequence, $q \in A$, therefore $A \in \mathcal{L}_{\beta}$.

(P4) Let $v \in T$ and $u \in \mathbf{Cl}(\mathcal{L}, \mathcal{L})$ such that $u \circ f = f \circ v$. Define $u_0 := F^{-1} \circ u \circ F$. Then $u_0 \in \text{Aut}(\mathcal{L}_0)$ and $u_0(p) = v(p)$ for all $p \in \Sigma_0$. $\square$

5. SUFFICIENT AND NECESSARY CONDITIONS FOR $\mathcal{L}_1 \otimes \mathcal{L}_2 = \mathcal{L}_1 \vee \mathcal{L}_2$

Let $\mathcal{L}_1, \dots, \mathcal{L}_n \in \mathbf{Cl}$ be DAC-lattices. In this section, we prove that $\bigotimes_{i=1}^n \mathcal{L}_i = \bigvee_{i=1}^n \mathcal{L}_i$ (i.e. $\mathbf{S}(\mathcal{L}_1, \dots, \mathcal{L}_n)$ has only one element) if and only if there is k between 1 and n such that for all $i \neq k$, $\mathcal{L}_i$ is a power set.

Lemma 5.1. *Let $\{\mathcal{L}_{\alpha} \in \mathbf{Cl}(\Sigma_{\alpha})\}_{\alpha \in \Omega}$ and $\mathcal{L} \in \mathbf{S}(\mathcal{L}_{\alpha}, \alpha \in \Omega)$. Let $\Sigma = \prod_{\alpha} \Sigma_{\alpha}$ and $R \subseteq \Sigma$. Define*

$$\vee_{\alpha} R := \bigcup_{p \in \Sigma} p[\vee R_{\alpha}[p]].$$

Denote by $\vee$ the join in $\mathcal{L}$. Then for any $f : \mathbb{N} \rightarrow \Omega$ and any $n \in \mathbb{N}$, $R^n := \vee_{f(n)} \dots \vee_{f(1)} R \subseteq \vee R$. Moreover, if $\vee_{\beta} R = R$ for all $\beta \in \Omega$, then $R \in \bigvee_{\alpha} \mathcal{L}_{\alpha}$.

Proof. Obviously, for all $\beta \in \Omega$, $\vee_{\beta}(\vee_{\beta} R) = \vee_{\beta} R$, and by Axioms P2 and P3, $\vee_{\beta} \vee R = \vee R$. Therefore, since $R \subseteq \vee R$, $R^n \subseteq (\vee R)^n = \vee R$.

Moreover, from Definition 2.11, $R \in \bigvee_{\alpha} \mathcal{L}_{\alpha}$ if and only if $\vee_{\beta} R = R$ for all $\beta \in \Omega$. $\square$

Theorem 5.2. *Let $\{\mathcal{L}_{\alpha} \in \mathbf{Cl}(\Sigma_{\alpha})\}_{\alpha \in \Omega}$ and $\Sigma = \prod_{\alpha} \Sigma_{\alpha}$. If there is at most one $\beta \in \Omega$ such that $\mathcal{L}_{\beta} \neq 2^{\Sigma_{\beta}}$, then $\bigotimes_{\alpha} \mathcal{L}_{\alpha} = \bigvee_{\alpha} \mathcal{L}_{\alpha} = \{R \subseteq \Sigma ; R_{\beta}[p] \in \mathcal{L}_{\beta}, \forall p \in \Sigma\}$.*

Proof. Let $R \subseteq \Sigma$. Since $\bigotimes_{\alpha} \mathcal{L}_{\alpha} \subseteq \bigvee_{\alpha} \mathcal{L}_{\alpha}$, $\vee_{\bigotimes} R \subseteq \vee_{\bigotimes} R$. We prove that $\vee_{\bigotimes} R \subseteq \vee_{\bigotimes} R$.

Write $\Sigma_1 = \Sigma_{\beta}$, $\mathcal{L}_1 = \mathcal{L}_{\beta}$, and $\Sigma_2 = \prod_{\alpha \neq \beta} \Sigma_{\alpha}$. Hence $\Sigma = \Sigma_1 \times \Sigma_2$. We denote by $\pi_i : \Sigma \rightarrow \Sigma_i$ ($i = 1, 2$) the i -th coordinate map. From definition 2.11, we find that

$$\bigotimes_{\alpha \neq \beta} \mathcal{L}_{\alpha} = 2^{\Sigma_2}.$$

Therefore, by Proposition 2.12,

$$\bigotimes_{\alpha \in \Omega} \mathcal{L}_{\alpha} = \mathcal{L}_1 \bigotimes 2^{\Sigma_2}.$$

Let $a \in \mathcal{L}_1$ and $b \in 2^{\Sigma_2}$. Note that $R \subseteq a \times \Sigma_2 \cup \Sigma_1 \times b$ if and only if

$$g(b) := \pi_1(\Sigma_1 \times b^c \cap R) \subseteq a,$$

where b^c denotes the set-complement of b , i.e. $b^c = \Sigma_2 \setminus b$. As a consequence, we find that

$$\begin{aligned} \vee_{\otimes} R &\subseteq \bigcap_{b \subseteq \pi_2(R)} (\vee g(b)) \times \Sigma_2 \cup \Sigma_1 \times b \\ &= \bigcup_{f \in 2^{(2^{\pi_2(R)})}} (\cap \{\vee g(b); b \in f^{-1}(1)\}) \times (\cap f^{-1}(2)). \end{aligned}$$

For $b \subseteq \pi_2(R)$, define $m(b) := \{a \subseteq \pi_2(R); b \not\subseteq a\}$. Then,

$$\vee_{\otimes} R = \bigcup_{b \subseteq \pi_2(R)} (\cap \{\vee g(a); a \in m(b)\}) \times b.$$

Note that for all $q \in b$, $q^c \cap \pi_2(R) \in m(b)$. Moreover,

$$g(q^c \cap \pi_2(R)) = \pi_1(\Sigma_1 \times (q \cup \pi_2(R)^c) \cap R) = R_1[(\cdot, q)].$$

As a consequence,

$$\begin{aligned} \vee_{\otimes} R &\subseteq \bigcup_{b \subseteq \pi_2(R)} (\cap \{\vee R_1[(\cdot, q)]; q \in b\}) \times b \\ &= \bigcup_{q \in \pi_2(R)} (\vee R_1[(\cdot, q)]) \times q = \vee_{\beta} R. \end{aligned}$$

Finally, by Lemma 5.1, $\vee_{\beta} R = \vee_{\otimes} R$. $\square$

Definition 5.3. Let Σ be a nonempty set. We denote by $\mathbf{MO}_{\Sigma}$ the simple closure space on Σ which contains only $\emptyset$, Σ , and all singletons of Σ . We write $\mathbf{MO}_n$ if $|\Sigma| = n$.

Let $\mathcal{L} \in \mathbf{Cl}$. We say that $\mathcal{L}$ contains $\mathbf{MO}_n$ if there are n atoms $p_1, \dots, p_n$ such that $p_1 \vee p_n \succ p_i$ for all i between 1 and n .

Let $\{\Sigma_{\alpha}\}_{\alpha \in \Omega}$ be a family of sets, and $\Sigma = \prod_{\alpha} \Sigma_{\alpha}$. We denote by $\Xi(\Sigma)$ the set $\{R \subseteq \Sigma; p_{\alpha} \neq q_{\alpha}, \forall p, q \in R, \alpha \in \Omega\}$.

Theorem 5.4. Let $\mathcal{L}_1 \in \mathbf{Cl}(\Sigma_1)$ and $\mathcal{L}_2 \in \mathbf{Cl}(\Sigma_2)$. If both $\mathcal{L}_1$ and $\mathcal{L}_2$ contain $\mathbf{MO}_3$, then $\mathcal{L}_1 \otimes \mathcal{L}_2 \neq \mathcal{L}_1 \vee \mathcal{L}_2$.

Proof. Let $R = \{p, q, r\} \in \Xi(\Sigma_1 \times \Sigma_2)$ such that for $i = 1$ and for $i = 2$, $p_i \vee q_i$ covers p_i, q_i and r_i . By Definition 2.11, $R = \{p, q, r\} \in \mathcal{L}_1 \vee \mathcal{L}_2$. On the other hand, $R \subseteq a \times \Sigma_2 \cup \Sigma_1 \times b$ if and only if $p_1 \vee q_1 \subseteq a$ or $p_2 \vee q_2 \subseteq b$. As a consequence,

$$\vee_{\otimes} R = (p_1 \vee q_1) \times (p_2 \vee q_2) \neq R = \vee_{\vee} R.$$

$\square$

Corollary 5.5. Let $\{\mathcal{L}_i \in \mathbf{Cl}(\Sigma_i)\}_{1 \leq i \leq n}$, be DAC-lattices. If $\bigotimes_{i=1}^n \mathcal{L}_i = \bigvee_{i=1}^n \mathcal{L}_i$, then there is at most one i such that $\mathcal{L}_i \neq 2^{\Sigma_i}$.

Proof. Suppose that $\mathcal{L}_i \neq 2^{\Sigma_i}$ for $i = m$ and for $i = k$ with $m \neq k$ between 1 and n , and that $\bigotimes_{i=1}^n \mathcal{L}_i = \bigvee_{i=1}^n \mathcal{L}_i$. Then, for $i = m$ and for $i = k$, there is an atom p_i which is not a central element. Therefore, there is an atom q_i such that $p_i \vee q_i$ contains a third atom (see [13], Theorems 28.8, 27.6 and Lemma 11.6).

Let $\mathcal{L}_0 := \bigotimes_{i \neq m} \mathcal{L}_i$ and $\mathcal{L}_1 := \bigotimes_{i \neq m} \mathcal{L}_i$. By hypothesis, $\mathcal{L}_m \bigotimes \mathcal{L}_0 = \mathcal{L}_m \bigotimes \mathcal{L}_1$. Therefore, from Axioms P2 and P3, $\mathcal{L}_0 = \mathcal{L}_1$. Let $r \in \prod_{i \neq m} \Sigma_i$. From Axiom P2, $r[p_k] \vee_{\bigotimes} r[q_k] = r[p_k \vee q_k]$, hence contains a third atom, therefore $\mathcal{L}_0$ contains MO_3 . As a consequence, from Theorem 5.4, we find that

$$\mathcal{L}_m \bigotimes \mathcal{L}_1 = \mathcal{L}_m \bigotimes \mathcal{L}_0 \neq \mathcal{L}_m \bigotimes \mathcal{L}_0 = \mathcal{L}_m \bigotimes \mathcal{L}_1,$$

a contradiction. $\square$

Theorem 5.6. *Let $\mathcal{L} \in \mathbf{Cl}(\Sigma)$ be coatomistic. Suppose that for all countable set A of coatoms $\mathcal{L}$, $\bigcup A \neq \Sigma$. For all integer i , let $\mathcal{L}_i = \mathcal{L}$. Then $\bigotimes_{i=1}^{\infty} \mathcal{L}_i \neq \bigotimes_{i=1}^{\infty} \mathcal{L}_i$.*

Proof. For all i , let $\Sigma_i = \Sigma$. Denote $\prod_{i=1}^{\infty} \Sigma_i$ by Σ . Let $R = \{p \in \Sigma; p_n = p_m, \forall m, n\} \neq \Sigma$. From Definition 2.11, $R \in \bigotimes_{i=1}^{\infty} \mathcal{L}_i$. On the other hand, by hypothesis, for all $a \in \prod_{i=1}^{\infty} \mathcal{L}_i$, $R \not\subseteq \bigcup_{i=1}^{\infty} \pi_i^{-1}(a_i)$, hence $\vee_{\bigotimes} R = 1$. $\square$

Example 5.7. The Hypothesis of Theorem 5.6 is fulfilled for instance if $\mathcal{L} = \mathbf{P}(\mathcal{H})$ with $\mathcal{H}$ a real or complex Hilbert space.

6. CENTRAL ELEMENTS

Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}\}_{\alpha \in \Omega}$, $\mathcal{L} \in \mathbf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$ and $\beta \in \Omega$. In this section we prove that if z is a central element of $\mathcal{L}_\beta$, then $\pi_\beta^{-1}(z)$ is a central element of $\mathcal{L}$. As a corollary, if $\mathcal{L}$ is irreducible, then all $\mathcal{L}_\alpha$'s are irreducible. We give some sufficient conditions under which the converse result holds for the bottom element $\bigotimes_\alpha \mathcal{L}_\alpha$, and we describe the decomposition into irreducible components of $\bigotimes_\alpha \mathcal{L}_\alpha$ in terms of the decompositions of each $\mathcal{L}_\alpha$.

Definition 6.1. Let a and b be elements of a lattice. Then (a, b) is said to be a *modular pair* (in symbols $(a, b)M$) if $(c \vee a) \wedge b = c \vee (a \wedge b)$ for all $c \leq b$.

Lemma 6.2. *Let $\mathcal{L} \in \mathbf{Cl}(\Sigma)$ and $z \in \mathcal{L}$. Then z is a central element of $\mathcal{L}$ if and only if $z^c := \Sigma \setminus z \in \mathcal{L}$ and $(z, z^c)M$ and $(z^c, z)M$.*

Proof. Direct from Theorem 4.13 (ε) in [13]. $\square$

Corollary 6.3. *Let $\mathcal{L} \in \mathbf{Cl}(\Sigma)$ be orthocomplemented and let $z \in \mathcal{L}$. Then z is a central element of $\mathcal{L}$ if and only if $z^\perp = z^c$.*

Proof. First, note that $a^\perp \subseteq a^c$ for all $a \in \mathcal{L}$. Suppose that z is a central element of $\mathcal{L}$. Then, since $(z, z^c)M$, we find that

$$z^c = (z^\perp \vee z) \cap z^c = z^\perp \vee (z \cap z^c) = z^\perp.$$

Suppose now that $z^\perp = z^c$. We prove that $(z, z^c)M$. The proof for $(z^c, z)M$ is similar. Let $c \in \mathcal{L}$ with $c \subseteq z^c$. Then $z = (z^c)^\perp \subseteq c^\perp$, therefore

$$(c \vee z) \cap z^c = (c \cup z)^\perp \cap z^\perp = [(c^\perp \cap z^c) \cup z]^\perp = c^{\perp\perp} = c.$$

$\square$

Theorem 6.4. *Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$, $\mathcal{L} \in \mathbf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$ and $\beta \in \Omega$. If z is a central element of $\mathcal{L}_\beta$, then $\pi_\beta^{-1}(z)$ is a central element of $\mathcal{L}$.*

Proof. From Lemma 6.2, $z^c := \Sigma_\beta \setminus z \in \mathcal{L}_\beta$. Therefore, by Axiom P2, $(\pi_\beta^{-1}(z))^c = \pi_\beta^{-1}(z^c) \in \mathcal{L}$. We denote $\pi_\beta^{-1}(z)$ by Z and $\pi_\beta^{-1}(z^c)$ by Z^c . We prove that $(Z, Z^c)M$. The proof for $(Z^c, Z)M$ is similar.

Let $R \in \mathcal{L}$ with $R \subseteq Z^c$. Let $\Sigma = \prod_\alpha \Sigma_\alpha$. Write R as $R = \cup\{p[R_\beta[p]] ; p \in \Sigma\}$. By Lemma 5.1, $R \vee Z = \cup\{p[R_\beta[p] \vee z] ; p \in \Sigma\}$. Therefore, since (z, z^c) is modular pair of $\mathcal{L}_\beta$, we find that

$$(R \vee Z) \cap Z^c = \bigcup_{p \in \Sigma} p[(R_\beta[p] \vee z) \cap z^c] = \bigcup_{p \in \Sigma} p[R_\beta[p]] = R.$$

□

Corollary 6.5. *Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}\}_{\alpha \in \Omega}$ and $\mathcal{L} \in \mathbf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$. If $\mathcal{L}$ is irreducible, then all $\mathcal{L}_\alpha$'s are irreducible.*

Definition 6.6. Let $\mathcal{L} \in \mathbf{Cl}(\Sigma)$, $p \in \Sigma$, and $a \in \mathcal{L}$. We denote the center of $\mathcal{L}$ by $\mathcal{Z}(\mathcal{L})$, we write $e(p) = \cap\{z \in \mathcal{Z}(\mathcal{L}) ; p \in z\}$ and $\mathcal{A}(\mathcal{Z}(\mathcal{L})) = \{e(p) ; p \in \Sigma\}$, and $[0, a]$ denotes the interval $\{b \in \mathcal{L} ; 0 \subseteq b \subseteq a\}$.

Lemma 6.7. *Let $\mathcal{L} \in \mathbf{Cl}(\Sigma)$ be orthocomplemented. Then $\mathcal{Z}(\mathcal{L})$ is a complete atomistic sublattice of $\mathcal{L}$, the set of atoms of which is given by $\mathcal{A}(\mathcal{Z}(\mathcal{L}))$. Moreover, $\Sigma = \coprod\{z ; z \in \mathcal{A}(\mathcal{Z}(\mathcal{L}))\}$, and*

$$\mathcal{L} \cong \prod_{z \in \mathcal{A}(\mathcal{Z}(\mathcal{L}))} [0, z],$$

ordered by $f \leq g \Leftrightarrow f(z) \leq g(z)$ for all $z \in \mathcal{A}(\mathcal{Z}(\mathcal{L}))$.

Proof. Let $\omega \subseteq \mathcal{Z}(\mathcal{L})$. Then

$$(\cap \omega)^c \supseteq (\cap \omega)^\perp = \left(\bigcap_{z \in \omega} z^{\perp\perp} \right)^\perp = \left(\bigcup_{z \in \omega} z^c \right)^{\perp\perp} = ((\cap \omega)^c)^{\perp\perp},$$

therefore, $\cap \omega \in \mathcal{Z}(\mathcal{L})$ by Corollary 6.3. Hence, $\mathcal{Z}$ is a complete sublattice of $\mathcal{L}$.

Let $p \in \Sigma$ and $r \in e(p)$. Then $e(r) \subseteq e(p)$. Suppose that p is not in $e(r)$. Then $p \in e(r)^c \cap e(p)$, hence $e(p) \subseteq e(r)^c \cap e(p)$, therefore $e(p) \subseteq e(r)^c$; whence $e(r) \subseteq e(p)^c \cap e(p)$, a contradiction. As a consequence, $p \in e(r)$, therefore $e(p) \subseteq e(r)$, hence $e(p) = e(r)$.

Let $p, q \in \Sigma$. We have proved that either $e(p) = e(q)$, or $e(p) \cap e(q) = 0$. As a consequence, $\Sigma = \coprod\{z ; z \in \mathcal{A}(\mathcal{Z}(\mathcal{L}))\}$, hence $\mathcal{A}(\mathcal{Z}(\mathcal{L}))$ is the set of atoms of $\mathcal{Z}(\mathcal{L})$, and $\mathcal{Z}(\mathcal{L})$ is atomistic. The rest of the proof follows directly from Lemma 5.1 in [13]. □

Theorem 6.8 ([12], Theorem 1). *Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$ with for all $\alpha \in \Omega$, $\mathcal{L}_\alpha$ orthocomplemented. Suppose that one of the following assumptions holds.*

- (1) Ω is finite.
- (2) For all $\alpha \in \Omega$, $\mathcal{L}_\alpha$ has the covering property, and for all $p \neq q \in \Sigma_\alpha$ with $e(p) = e(q)$, $p \vee q$ contains an infinite number of atoms.

Then, $\bigotimes_\alpha \mathcal{L}_\alpha$ is irreducible if and if all $\mathcal{L}_\alpha$'s are irreducible. Moreover,

$$\mathcal{A}(\mathcal{Z}(\bigotimes_\alpha \mathcal{L}_\alpha)) = \left\{ \prod_\alpha Z_\alpha ; Z \in \prod_\alpha \mathcal{A}(\mathcal{Z}(\mathcal{L}_\alpha)) \right\}.$$

7. AUTOMORPHISMS

Let $\mathcal{L}_1, \dots, \mathcal{L}_n \in \mathbf{Cl}$ different from 2, $\mathcal{L} \in \mathbf{S}(\mathcal{L}_1, \dots, \mathcal{L}_n)$ and $u \in \mathbf{Cl}(\mathcal{L}, \mathcal{L})$ sending atoms to atoms. In this section we prove that if u is *large* (see Definition 7.3 below), then there is a permutation f and arrows $v_i \in \mathbf{Cl}(\mathcal{L}_i, \mathcal{L}_{f(i)})$ sending atoms to atoms such that for any atom p of $\mathcal{L}$, $u(p)_{f(i)} = v_i(p_i)$. We need some hypotheses on each $\mathcal{L}_i$ which are true for instance if each $\mathcal{L}_i$ is irreducible orthocomplemented with the covering property or an irreducible DAC-lattice. Note that our hypotheses imply irreducibility. Note also that if u is an automorphism, then u is large.

Definition 7.1. Let $\mathcal{L} \in \mathbf{Cl}(\Sigma)$. We say that $\mathcal{L}$ is *weakly connected* if $\mathcal{L} \neq 2$ and if there is a *connected covering* of Σ , that is a family of subsets $\{A^\gamma \subseteq \Sigma; \gamma \in \sigma\}$ such that

- (1) $\Sigma = \cup\{A^\gamma; \gamma \in \sigma\}$ and $|A^\gamma| \geq 2$ for all $\gamma \in \sigma$,
- (2) for all $\gamma \in \sigma$ and all $p \neq q \in A^\gamma$, $p \vee q$ contains a third atom,
- (3) for all $p, q \in \Sigma$, there is a finite subset $\{\gamma_1, \dots, \gamma_n\} \subseteq \sigma$ such that $p \in A^{\gamma_1}$, $q \in A^{\gamma_n}$, and such that $|A^{\gamma_i} \cap A^{\gamma_{i+1}}| \geq 2$ for all $1 \leq i \leq n-1$.

We say that $\mathcal{L}$ is *connected* if $\mathcal{L} \neq 2$ and for all $p, q \in \Sigma$, $p \vee q$ contains a third atom, say r , such that $p \in q \vee r$ and $q \in p \vee r$.

Remark 7.2. Note that in part 2 of Definition 7.1, it is not required that the third atom under $p \vee q$ is in A^γ . Let $\mathcal{L} \in \mathbf{Cl}(\Sigma)$. If $\mathcal{L}$ is weakly connected, then $\mathcal{L}$ is irreducible (follows directly from Lemma 6.2 and Lemma 5.1 in [13], Theorem 4.13). On the other hand, if $\mathcal{L} \neq 2$ and $\mathcal{L}$ is irreducible orthocomplemented with the covering property or an irreducible DAC-lattice, then $\mathcal{L}$ is connected.

Definition 7.3. Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$, with for all $\alpha \in \Omega$, $\mathcal{L}_\alpha \neq 2$. Let $\mathcal{L} \in \mathbf{S}(\mathcal{L}_\alpha, \alpha \in \Omega)$, $\Sigma = \prod_{\alpha} \Sigma_\alpha$, and let $u \in \mathbf{Cl}(\mathcal{L}, \mathcal{L})$. We say that u is *large* if for all $\beta \in \Omega$ and $p \in \Sigma$, $u(p[\Sigma_\beta])$ is not an atom and $u(1) \not\subseteq \pi_\beta^{-1}(p_\beta)$.

Lemma 7.4. Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$ and $\mathcal{L} \in \mathbf{Cl}(\Sigma)$ with $\Sigma = \prod_{\alpha} \Sigma_\alpha$. Suppose that Axiom P2 holds in $\mathcal{L}$. Let $p, q \in \Sigma$.

- (1) If $p_\beta \neq q_\beta$ for at least two $\beta \in \Omega$, then $p \vee q = p \cup q$.
- (2) For all $\beta \neq \gamma \in \Omega$ and for all $b \in \mathcal{L}_\beta$ and $c \in \mathcal{L}_\gamma$ such that $p_\beta \in b$ and $p_\gamma \in c$, $p[b, \beta] \vee p[c, \gamma] = p[b, \beta] \cup p[c, \gamma]$.

Proof. (1) Define $\Omega_\neq := \{\beta \in \Omega; p_\beta \neq q_\beta\}$. Let $r \in p \vee q$, and $\beta \neq \gamma \in \Omega_\neq$. Then

$$\begin{aligned} p, q &\in [\pi_\beta^{-1}(p_\beta) \cup \pi_\gamma^{-1}(q_\gamma)] \cap [\pi_\beta^{-1}(q_\beta) \cup \pi_\gamma^{-1}(p_\gamma)] \\ &= [\pi_\beta^{-1}(p_\beta) \cap \pi_\gamma^{-1}(p_\gamma)] \cup [\pi_\beta^{-1}(q_\beta) \cap \pi_\gamma^{-1}(q_\gamma)]. \end{aligned}$$

Therefore, $r_\beta = x_\beta$ and $r_\gamma = x_\gamma$, with $x = p$ or q .

(2) We have

$$\cap_{\alpha \neq \beta, \gamma} \pi_\alpha^{-1}(b) \cap \pi_\alpha^{-1}(c) \cap [\pi_\beta^{-1}(p_\beta) \cup \pi_\gamma^{-1}(p_\gamma)] \cap \pi_\alpha^{-1}(p_\alpha) = p[c, \gamma] \cup p[b, \beta].$$

□

Theorem 7.5. Let $\{\mathcal{L}_i \in \mathbf{Cl}(\Sigma_i)\}_{i \in \Omega}$ with Ω finite such that for all $i \in \Omega$, $\mathcal{L}_i$ is connected. Let $\mathcal{L} \in \mathbf{S}(\mathcal{L}_i, i \in \Omega)$ and $u \in \mathbf{Cl}(\mathcal{L}, \mathcal{L})$ large, sending atoms to atoms. Then there is a bijection f of Ω , and for each $i \in \Omega$, there is $v_i \in \mathbf{Cl}(\mathcal{L}_i, \mathcal{L}_{f(i)})$ sending atoms to atoms such that $u(p)_{f(i)} = v_i(p_i)$ for all $p \in \Sigma$ and $i \in \Omega$.

Proof. The proof is similar to the proof of Theorem 3 in [12].

(1) Let $p \in \Sigma$ and $j \in \Omega$. **Claim:** There is $k \in \Omega$ such that

$$u(p[\Sigma_j]) \subseteq u(p)[\Sigma_k].$$

[*Proof:* Since $\mathcal{L}_j$ is connected, for all $q_j \in \Sigma_j$ different from p_j , $p_j \vee q_j$ contains a third atom, say r_j , and $p_j \in q_j \vee r_j$ and $q_j \in p_j \vee r_j$. Suppose that $u(p)_k \neq u(p[q_j])_k$ for at least two indices k . Then, by Lemma 7.4 part 1 and Axiom P3,

$$u(p[r_j]) \in u(p \vee p[q_j]) = u(p) \vee u(p[q_j]) = u(p) \cup u(p[q_j]).$$

Assume for instance that $u(p[r_j]) = u(p)$. Then,

$$u(p[q_j]) \in u(p \vee p[r_j]) = u(p) \vee u(p[r_j]) = u(p),$$

a contradiction.]

(2) Let $j \in \Omega$. Denote the k of part 1 by $f(j, p)$. **Claim:** The map $p \mapsto f(j, p)$ is constant. [*Proof:* Let $p, q \in \Sigma$ that differ only by one component, say $j' \neq j$, i.e. $p_{j'} \neq q_{j'}$, and $p_i = q_i$, for all $i \neq j'$. Suppose that $f(j, p) \neq f(j, q)$. Write $k := f(j, p)$ and $k' := f(j, q)$.

(2.1) We first prove that

$$u(p[\Sigma_j]) \vee u(q[\Sigma_j]) = u(p[\Sigma_j]) \cup u(q[\Sigma_j]).$$

By hypothesis,

$$u(p[\Sigma_j]) \subseteq u(p)[\Sigma_k] \text{ and } u(q[\Sigma_j]) \subseteq u(q)[\Sigma_{k'}].$$

Hence, for all $r_j \in \Sigma_j$, we have

$$u(p)_l = u(p[r_j])_l \forall l \neq k \text{ and } u(q)_m = u(q[r_j])_m \forall m \neq k'.$$

Since u is large, $u(p[\Sigma_j])$ is not an atom, hence there is $r_j \in \Sigma_j$ such that

$$u(p[r_j])_k \neq u(q)_k.$$

Note that $u(q)_k = u(q[r_j])_k$. Therefore, since $p[r_j]$ and $q[r_j]$ differ only by one component, by part 1 we have

$$u(p[r_j])_l = u(q[r_j])_l \forall l \neq k.$$

As a consequence,

$$u(q)_l = u(q[r_j])_l = u(p[r_j])_l = u(p)_l \forall l \neq k, k',$$

and

$$u(p)_{k'} = u(p[r_j])_{k'} = u(q[r_j])_{k'} \in \pi_{k'}(u(q[\Sigma_j])).$$

On the other hand, since $u(q[\Sigma_j])$ is not an atom, there is $s_j \in \Sigma_j$ such that

$$u(q[s_j])_{k'} \neq u(p)_{k'}.$$

Note that $u(p)_{k'} = u(p[r_j])_{k'}$. Therefore, since $q[s_j]$ and $p[s_j]$ differ only by one component, by part 1 we have

$$u(q[s_j])_l = u(p[s_j])_l \forall l \neq k'.$$

As a consequence,

$$u(q)_k = u(q[s_j])_k = u(p[s_j])_k \in \pi_k(u(p[\Sigma_j])).$$

To summarize, we have proved that $u(p)_l = u(q)_l$ for all $l \neq k, k'$, $u(p)_{k'} \in \pi_{k'}(u(q[\Sigma_j]))$, and $u(q)_k \in \pi_k(u(p[\Sigma_j]))$. As a consequence, the statement follows from Lemma 7.4 part 2.

(2.2) Since $\mathcal{L}_{j'}$ is connected, there is $s_{j'} \in p_{j'} \vee q_{j'}$ such that $p_{j'} \in q_{j'} \vee s_{j'}$ and $q_{j'} \in p_{j'} \vee s_{j'}$. Let $r = p[s_{j'}]$. Then, by Axiom P3, for all $t_j \in \Sigma_j$, $r[t_j] \subseteq p[t_j] \vee q[t_j]$. Therefore

$$r[\Sigma_j] \subseteq p[\Sigma_j] \vee q[\Sigma_j],$$

and for the same reason

$$p[\Sigma_j] \subseteq q[\Sigma_j] \vee r[\Sigma_j] \text{ and } q[\Sigma_j] \subseteq p[\Sigma_j] \vee r[\Sigma_j].$$

Now, by part 2.1,

$$u(p[\Sigma_j] \vee q[\Sigma_j]) = u(p[\Sigma_j]) \vee u(q[\Sigma_j]) = u(p[\Sigma_j]) \cup u(q[\Sigma_j]).$$

As a consequence, by part 1, $u(r[\Sigma_j]) \subseteq u(p[\Sigma_j])$ or $u(r[\Sigma_j]) \subseteq u(q[\Sigma_j])$. Assume for instance that $u(r[\Sigma_j]) \subseteq u(p[\Sigma_j])$. Then,

$$u(q[\Sigma_j]) \subseteq u(p[\Sigma_j] \vee r[\Sigma_j]) = u(p[\Sigma_j]) \vee u(r[\Sigma_j]) = u(p[\Sigma_j]),$$

a contradiction. Hence, we have proved that if $p, q \in \Sigma$ differ only by one component, then $f(j, p) = f(j, q)$.

(2.3) Suppose now that p and q differ by more than one component. Since Ω is finite, there is $s_1, \dots, s_n \in \Sigma$ such that $s^1 = q$, $s^n = p$, and such that for all $1 \leq i \leq n-1$, s^i and s^{i+1} differ only by one component. Therefore,

$$f(j, q) = f(j, s^1) = f(j, s^2) = \dots = f(j, s^n) = f(j, p),$$

and we are done.]

(3) Let $p_0 \in \Sigma$. Define $f : \Omega \rightarrow \Omega$ as $f(i) := f(i, p_0)$. Note that by part 2, f does not depend on the choice of p_0 . **Claim:** The map f is surjective. [*Proof:* Let $k \in \Omega$. Suppose that for all $i \in \Omega$ $f(i) \neq k$. Let $p, q \in \Sigma$ that differ only by one component, say j . Then $p[\Sigma_j] = q[\Sigma_j]$, therefore $u(p[\Sigma_j]) = u(q[\Sigma_j])$. Moreover, by part 1, there is $k' \neq k$ such $u(p[\Sigma_j]) \subseteq u(p)[\Sigma_{k'}]$. As a consequence, $u(p)_k = u(q)_k$.

Since Ω is finite, by the same argument as in part 2.3, we find that $u(p)_k = u(q)_k$, for all $p, q \in \Sigma$. As a consequence, $u(\Sigma) \subseteq \pi_k^{-1}(u(p_0)_k)$, a contradiction since u is large.]

(4) Let $p_0 \in \Sigma$ and $j \in \Omega$. Define $v_j : \mathcal{L}_j \rightarrow \mathcal{L}_{f(j)}$ as $v_j(a_j) := \pi_{f(j)}(u(p_0[a_j]))$. **Claim:** v_j does not depend on the choice of p_0 . [*Proof:* Let $q \in \Sigma$ that differs from p_0 only by one component, say $j' \neq j$. Then, by Axiom P3, we have

$$\begin{aligned} \pi_{f(j)}(u(q[a_j])) &= \pi_{f(j)}(u(\vee\{q[r_j]; r_j \in a_j\})) \\ &= \pi_{f(j)}(\vee\{u(q[r_j]); r_j \in a_j\}) = \bigvee_{r_j \in a_j} u(q[r_j])_{f(j)}, \end{aligned}$$

and the same formula holds for $\pi_{f(j)}(u(p_0[a_j]))$. Now

$$u(q[r_j])_{f(j)} = \pi_{f(j)}(u(s[\Sigma_{j'}])) = u(p_0[r_j])_{f(j)},$$

where $s = q[r_j]$. As a consequence, $\pi_{f(j)}(u(q[a_j])) = \pi_{f(j)}(u(p_0[a_j]))$.

Since Ω is finite, by the same argument as in part 2.3, we find that $\pi_{f(j)}(u(q[a_j])) = \pi_{f(j)}(u(p_0[a_j]))$ for all $q \in \Sigma$.]

It remains to check that v_j preserves arbitrary joins. Let $\omega \subseteq \mathcal{L}_j$. Then, by Axiom P3,

$$\begin{aligned} v_j(\vee\omega) &= \pi_{f(j)}(u(p[\vee\omega])) = \pi_{f(j)}(u(\vee\{p[x, j]; x \in \omega\})) \\ &= \pi_{f(j)}(\vee\{u(p[x, j]); x \in \omega\}) \\ &= \bigvee_{x \in \omega} \pi_{f(j)}(u(p[x, j])) = \bigvee_{x \in \omega} v_j(x). \end{aligned}$$

□

Corollary 7.6. *If the u in Theorem 7.5 is an automorphism, then all v_i 's are isomorphisms.*

Theorem 7.7. *If the u in Theorem 7.5 is an automorphism, the statement remains true if for all $i \in \Omega$, $\mathcal{L}_i$ is weakly connected.*

Proof. The proof is similar as in Theorem 7.5. We only sketch the arguments that must be modified. Note that since u is an automorphism and $\mathcal{L}_i \neq 2$ for all $i \in \Omega$, u is large.

(1) Since $\mathcal{L}_j$ is weakly connected, there is $\gamma_0 \in \sigma^j$ such that $p_j \in A_j^{\gamma_0}$. By hypothesis and Axiom P3, for all $q_j \in A_j^{\gamma_0}$, $p \vee (p[q_j])$ contains a third atom, hence also $u(p) \vee u(p[q_j])$ since u is injective. As a consequence, there is $k_{\gamma_0} \in \Omega$ such that $u(p[A_j^{\gamma_0}]) \subseteq u(p)[\Sigma_{k_{\gamma_0}}]$. Moreover, since u is injective, by the third hypothesis in Definition 7.1, the map $\gamma \mapsto k_\gamma$ is constant. Therefore, since $\cup A_j^\gamma = \Sigma_j$, we find that $u(p[\Sigma_j]) \subseteq u(p)[\Sigma_k]$.

(2) Take $p, q \in \Sigma$ that differ only by one component such that $q_{j'}$ and $p_{j'}$ are in the same A_j^γ .

(2.2) By hypothesis, $p_{j'} \vee q_{j'}$ contains a third atom, say $r_{j'}$, therefore $r[\Sigma_j] \subseteq p[\Sigma_j] \vee q[\Sigma_j]$, hence, $u(r[\Sigma_j]) = u(p[\Sigma_j])$ or $u(r[\Sigma_j]) = u(q[\Sigma_j])$, a contradiction since u is injective. As a consequence, $f(j, q) = f(j, p)$. Now, since u is injective, by the third hypothesis in definition 7.1, we find that $f(j, q) = f(j, p)$, for all $p, q \in \Sigma$ that differ only by the component j' . □

8. ORTHOCOMPLEMENTATION

This section is devoted to our main result, namely that if $\mathcal{L} \in \mathbf{S}(\mathcal{L}_1, \dots, \mathcal{L}_n)$ is orthocomplemented, then $\mathcal{L} = \bigotimes_{i=1}^n \mathcal{L}_i$. We provide two proofs. The first one uses Theorem 7.7, hence requires each $\mathcal{L}_i$ to be weakly connected, so in particular irreducible. In addition we must assume that each $\mathcal{L}_i$ is coatomistic and that $\mathcal{L}$ is transitive. For the second proof, we assume that each $\mathcal{L}_i$ is orthocomplemented and that all its irreducible components different from 2 are weakly connected, but we do not need to assume that $\mathcal{L}$ is transitive.

Definition 8.1. Let $\{\Sigma_\alpha\}_{\alpha \in \Omega}$ be a family of nonempty sets, $p \in \Sigma = \prod_{\alpha} \Sigma_\alpha$, $\beta, \gamma \in \Omega$, $A_\beta \subseteq \Sigma_\beta$, and $A_\gamma \subseteq \Sigma_\gamma$. Then

$$p[A_\beta, A_\gamma] := \{q \in \Sigma; q_\beta \in A_\beta, q_\gamma \in A_\gamma, q_\alpha = p_\alpha, \forall \alpha \neq \beta, \gamma\}.$$

Lemma 8.2. *Let $\{\mathcal{L}_i \in \mathbf{Cl}(\Sigma_i)\}_{1 \leq i \leq n}$, $x \in \prod_{i=1}^n \mathcal{L}_i$ with x_i coatoms, and let $\mathcal{L} \in \mathbf{S}(\mathcal{L}_i, 1 \leq i \leq n)$. Then $X := \cup_{i=1}^n \pi_i^{-1}(x_i)$ is a coatom of $\mathcal{L}$.*

Proof. By Axiom P2, $X \in \mathcal{L}$. Let $\Sigma = \prod_{i=1}^n \Sigma_i$ and $p \in \Sigma$ not in X . Write $R^0 := p \cup X$ and $y := p \vee X$. Define $R^N := \vee_n \dots \vee_1 R^{N-1}$. By Lemma 5.1, $R^N \subseteq y$, for all N . Now, $\vee_k R^0 = p[\Sigma_k] \cup R^0$, thus $R^1 = R^0 \cup_{k=1}^n p[\Sigma_k]$. Further,

$$R^2 = R^0 \bigcup_{k_1=1}^{n-1} \bigcup_{k_2=k_1+1}^n p[\Sigma_{k_1}, \Sigma_{k_2}].$$

Hence obviously, for $N = n$, $R^N = \Sigma$. □

Proposition 8.3. *Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$.*

- (1) If $\bigotimes_{\alpha} \mathcal{L}_{\alpha}$ is coatomistic, then all $\mathcal{L}_{\alpha}$'s are coatomistic.
 (2) Suppose that Ω is finite and that all $\mathcal{L}_{\alpha}$'s are weakly connected. If $\bigotimes_{\alpha} \mathcal{L}_{\alpha}$ is orthocomplemented, then all $\mathcal{L}_{\alpha}$'s are orthocomplemented.

Proof. (1) Let $\Sigma = \prod_{\alpha} \Sigma_{\alpha}$ and $p \in \Sigma$. From Definition 2.11, a coatom of $\bigotimes_{\alpha} \mathcal{L}_{\alpha}$ over p is necessarily of the form $\cup_{\alpha} \pi_{\alpha}^{-1}(x_{\alpha})$ where all x_{α} 's are coatoms and with $p_{\beta} \in x_{\beta}$ for some $\beta \in \Omega$. Let $\Sigma'[p_{\alpha}]$ denote the set of coatoms of $\mathcal{L}_{\alpha}$ above p_{α} (write $\Sigma'_{\alpha} := \Sigma'_{\alpha}[0_{\alpha}]$). If $\bigotimes_{\alpha} \mathcal{L}_{\alpha}$ is coatomistic, then

$$p = \bigcap \{ \cup_{\alpha} \pi_{\alpha}^{-1}(x_{\alpha}) ; x \in \prod_{\alpha} \Sigma'_{\alpha} \text{ with } p_{\beta} \in x_{\beta} \text{ for some } \beta \in \Omega \}.$$

As a consequence, all $\mathcal{L}_{\alpha}$'s are coatomic. Moreover, for all $\alpha \in \Omega$, $p_{\alpha} = \cap \{x_{\alpha} ; x_{\alpha} \in \Sigma'[p_{\alpha}]\}$, hence $\mathcal{L}_{\alpha}$ is coatomistic.

(2) Let $\beta \in \Omega$ and $a \in \mathcal{L}_{\beta}$. We first prove that there is $b \in \mathcal{L}_{\beta}$ such that $\pi_{\beta}^{-1}(a)' = \pi_{\beta}^{-1}(b)$. Let $f : \Sigma \rightarrow \prod \Sigma'_{\alpha}$ such that $p' = \cup_{\alpha} \pi_{\alpha}^{-1}(f_{\alpha}(p))$ for all $p \in \Sigma$.

(2.1) Let $p, q \in \Sigma$. **Claim:** If $f(p)_{\alpha} \neq f(q)_{\alpha}$ for at least two $\alpha \in \Omega$, then $p \vee q = p \cup q$. [*Proof:* Let $\Omega_{\neq} = \{\alpha \in \Omega ; f(p)_{\alpha} \neq f(q)_{\alpha}\}$ and $\gamma \neq \delta \in \Omega_{\neq}$. Let $r \in p \vee q$. Then,

$$\cup_{\alpha} \pi_{\alpha}^{-1}(f(r)_{\alpha}) = r' \supseteq p' \cap q' \supseteq \pi_{\gamma}^{-1}(f(p)_{\gamma}) \cap \pi_{\delta}^{-1}(f(q)_{\delta}) \cup \pi_{\gamma}^{-1}(f(q)_{\gamma}) \cap \pi_{\delta}^{-1}(f(p)_{\delta}).$$

Therefore, $f(r)_{\gamma} = x_{\gamma}$ and $f(r)_{\delta} = x_{\delta}$ with $x = f(p)$ or $x = f(q)$. As a consequence, $r' = p'$ or $r' = q'$, hence $r = p$ or $r = q$.]

(2.2) **Claim:** Let $p, q \in \Sigma$ such that p and q differ only by one component, say γ . Then $f(p)_{\alpha} = f(q)_{\alpha}$ for all $\alpha \neq \gamma$. [*Proof:* Since $\mathcal{L}_{\gamma}$ is weakly connected, from part 2.1 we find that there is $\delta \in \Omega$ such that for all $r \in \Sigma_{\gamma}$, $f(p[r, \gamma])_{\alpha} = f(p)_{\alpha}$ for all $\alpha \neq \delta$. Hence,

$$p[\Sigma_{\gamma}]' = \bigcup_{\alpha \neq \delta} \pi_{\alpha}^{-1}(f(p)_{\alpha}) \cup \pi_{\delta}^{-1}(x),$$

for some $x \in \mathcal{L}_{\delta}$. Now $p[\Sigma_{\gamma}] \cap \pi_{\gamma}^{-1}(f(p)_{\gamma}) \neq 0$. Therefore, if $\delta \neq \gamma$, then $p[\Sigma_{\gamma}] \cap p[\Sigma_{\gamma}]' \neq 0$, a contradiction. As a consequence, $\delta = \gamma$ and $x = 0$.]

Let $p, q \in \Sigma$ such that $p \in \pi_{\beta}^{-1}(a)'$ and such that p and q differ only by one component, say γ , different from β . Then $\pi_{\beta}^{-1}(a) \subseteq p' = \cup_{\alpha} \pi_{\alpha}^{-1}(f(p)_{\alpha})$, hence $a \subseteq f(p)_{\beta}$. Now, from part 2.2, $f(q)_{\beta} = f(p)_{\beta}$, therefore $\pi_{\beta}^{-1}(a) \subseteq q'$, thus $q \in \pi_{\beta}^{-1}(a)'$. Now, by the same argument as in the proof of Theorem 7.5 part 2.3, we find that for all $q \in \Sigma$ such that $q_{\beta} = p_{\beta}$, $q \in \pi_{\beta}^{-1}(a)'$.

As a consequence, we have proved that there is an element in $\mathcal{L}_{\beta}$, which we denote by $a^{\perp \beta}$, such that $\pi_{\beta}^{-1}(a)' = \pi_{\beta}^{-1}(a^{\perp \beta})$. Obviously, the mapping $\perp_{\beta} : \mathcal{L}_{\beta} \rightarrow \mathcal{L}_{\beta}$ is an orthocomplementation. $\square$

Theorem 8.4. Let $\{\mathcal{L}_i \in \mathbf{Cl}(\Sigma_i)\}_{1 \leq i \leq n}$, with for all i , $\mathcal{L}_i$ coatomistic and weakly connected, and let $\mathcal{L} \in \mathbf{S}(\mathcal{L}_i, 1 \leq i \leq n)$. If $\mathcal{L}$ is transitive and orthocomplemented, then $\mathcal{L} = \bigotimes_{i=1}^n \mathcal{L}_i$.

Proof. Let $\Sigma = \prod_{i=1}^n \Sigma_i$. We denote the orthocomplementation of $\mathcal{L}$ by $'$. Let $x \in \prod_{i=1}^n \mathcal{L}_i$ with x_i coatoms, and $X := \cup_{i=1}^n \pi_i^{-1}(x_i)$. By Lemma 8.2, $X' = p$, for some $p \in \Sigma$. Let $q \in \Sigma$. Since $\mathcal{L}$ is transitive, there is $u \in \mathbf{Aut}(\mathcal{L})$ such that $u(p) = q$, hence $q' = u(p)'$. Define $u' \in \mathbf{Aut}(\mathcal{L})$ as $u'(a) := u(a)'$. Then $q' = u(p)' = u'(p') = u'(X)$. From Theorem 7.7, u' factors, therefore there is $y \in \prod_{i=1}^n \mathcal{L}_i$ with y_i coatoms such that $q' = Y := \cup_{i=1}^n \pi_i^{-1}(y_i)$. $\square$

Remark 8.5. A similar result, using a different set of axioms than Axioms P1-P3, is established in [10] for $n = 2$ and $\mathcal{L}_i = \mathbf{P}(\mathcal{H}_i)$ with $\mathcal{H}_i$ complex Hilbert spaces. If $\mathcal{L} \in \mathcal{S}_T(\mathcal{L}_i, 1 \leq i \leq n)$, $\mathcal{L}_i$ are transitive, and the u in Axiom P4 is an ortho-isomorphism of $\mathcal{L}$ for all $v_i \in T_i$, then the proof does not require Theorem 7.7. Below we give a second proof which requires neither Theorem 7.7 nor $\mathcal{L}$ to be transitive.

Theorem 8.6. *Let $\{\mathcal{L}_i \in \mathbf{Cl}(\Sigma_i)\}_{1 \leq i \leq n}$, with for all i , $\mathcal{L}_i$ orthocomplemented. Suppose moreover that for all i , all irreducible components of $\mathcal{L}_i$ different from 2 are weakly connected. Let $\mathcal{L} \in \mathbf{S}(\mathcal{L}_i, 1 \leq i \leq n)$. If $\mathcal{L}$ is orthocomplemented, then $\mathcal{L} = \bigotimes_{i=1}^n \mathcal{L}_i$.*

Proof. Let $\Sigma = \prod_{i=1}^n \Sigma_i$. We denote the orthocomplementation of $\mathcal{L}$ by $'$. From Lemma 8.2, we can define a map ϕ on Σ as $\phi(p) = p^{\#}'$ (see Definition 3.1). Note that ϕ is injective. We prove in four steps that ϕ is surjective.

(1) Let $p \in \Sigma$ and $a_j \in \mathcal{L}_j$ for some j between 1 and n . **Claim:** $\phi(p[a_j]) = p[a_j]^{\#}'$. [*Proof:* If $q \in p[a_j]$, then $p[a_j]^{\#} \subseteq q^{\#}$, hence $\phi(q) \in p[a_j]^{\#}'$. On the other hand, if $q \in p[a_j]^{\#}'$, then,

$$p[a_j]^{\#} = \bigcup_{i \neq j} \pi_i^{-1}(p_i^{\perp i}) \cup \pi_j^{-1}(a_j^{\perp j}) \subseteq q'.$$

Let $r \in q'$ not in $p[a_j]^{\#}$. Note that $r[a_j^{\perp j}] \subseteq p[a_j]^{\#}$. Therefore, by Lemma 5.1, $r[a_j^{\perp j} \vee r_j] \subseteq q'$. Let $k \neq j$ between 1 and n . Then, $r[p_k^{\perp k}, a_j^{\perp j} \vee r_j] \subseteq p[a_j]^{\#}$. Hence, since $r[a_j^{\perp j} \vee r_j] \subseteq q'$, by Lemma 5.1, $r[\Sigma_k, a_j^{\perp j} \vee r_j] \subseteq q'$. Repeating this argument as in the proof of Lemma 8.2, we find that $\pi_j^{-1}(a_j^{\perp j} \vee r_j) \subseteq q'$. As a consequence, there is an atom s_j with $a_j^{\perp j} \subseteq s_j^{\perp j}$ such that

$$q' = \bigcup_{i \neq j} \pi_i^{-1}(p_i^{\perp i}) \cup \pi_j^{-1}(s_j^{\perp j}).$$

Therefore, $q = p[s_j]^{\#}'$ with $s_j \in a_j$, hence $q \in \phi(p[a_j])$.

(2) Let $p \in \Sigma$. **Claim:** For all j between 1 and n there is $q \in \prod_{i=1}^n e(p_i)$ and k between 1 and n such that $\phi(p[e(p_j)]) \subseteq q[e(p_k)]$ (see Definition 6.6). [*Proof:* Note that if $a \in \mathcal{L}$, then $a' \subseteq a^c$, where a^c denotes the set-complement of a , i.e. $a^c = \Sigma \setminus a$. By part 1,

$$\phi(p[e(p_j)]) \subseteq (p[e(p_j)]^{\#})^c = e(p_j) \prod_{i \neq j} ((p_i^{\perp i})^c) \subseteq \prod_{i=1}^n e(p_i),$$

since if $q_i \in e(p_i)$, then $(q_i^{\perp i})^c \subseteq e(p_i)$. If $[0, e(p_j)] = 2$ (i.e. $e(p_j) = p_j$), then the proof is finished.

Otherwise, let $t_j \neq s_j \in e(p_j)$. Then,

$$\begin{aligned} q \in \phi(p[t_j]) \vee \phi(p[s_j]) &\Leftrightarrow p[t_j]^{\#} \cap p[s_j]^{\#} \subseteq q' \\ &\Leftrightarrow \bigcup_{i \neq j} \pi_i^{-1}(p_i^{\perp i}) \cup \pi_j^{-1}((t_j \vee s_j)^{\perp j}) \subseteq q', \end{aligned}$$

hence if and only if $q = \phi(p[r_j])$ for some $r_j \in t_j \vee s_j$. Therefore, if $t_j \vee s_j$ contains a third atom, so does $\phi(p[t_j]) \vee \phi(p[s_j])$. Hence, by Lemma 7.4 part 1, $\phi(p[t_j])$ and $\phi(p[s_j])$ differ only by one component, say k . As a consequence, since $\mathcal{L}_j$ is weakly connected, for all $A_j^?$ in the connected covering of Σ_j , there is $q^? \in \prod_{i=1}^n e(p_i)$ and

k_γ such that $\phi(p[e(p_j) \cap A_j^\gamma]) \subseteq q^\gamma[e(p_{k_\gamma})]$. From Hypotheses 1 and 3 in Definition 7.1, the maps $\gamma \mapsto k_\gamma$ and $\gamma \mapsto q_\gamma$ are constant since ϕ is injective.]

(3) **Claim:** For all $p \in \Sigma$ and all j between 1 and n , there is $q \in \prod_{i=1}^n e(p_i)$ such that $\phi(p[e(p_j)]) = q[e(p_j)]$. [*Proof:* From part 2, there is $q \in \prod_{i=1}^n e(p_i)$, k between 1 and n , and $b_k \subseteq e(p_k)$, such that $\phi(p[e(p_j)]) = q[b_k]$.

Assume first that $k \neq j$. Let $R^0 := q[b_k] \cup p[e(p_j)]^\#$. By part 1,

$$R^{0'} = q[b_k]' \cap p[e(p_j)]^{\#'} = \phi(p[e(p_j)])' \cap \phi(p[e(p_j)]) = 0,$$

hence $\vee R^0 = 1$. On the other hand,

$$R^0 = q[b_k] \bigcup_{i \neq j} \pi_i^{-1}(p_i^{\perp i}) \cup \pi_j^{-1}(e(p_j)^c) \subseteq \pi_j^{-1}(r_j^{\perp j}) \bigcup_{i \neq j} \pi_i^{-1}(p_i^{\perp i}),$$

for all $r_j \in e(p_j) \cap q_j^{\perp j}$, whence by Axiom P2, $\vee R^0 \neq 1$, a contradiction. As a consequence, $k = j$.

Let $R := q[b_j] \cup p[e(p_j)]^\#$. From part 1, $R' = \phi(p[e(p_j)])' \cap \phi(p[e(p_j)]) = 0$. Therefore, $\vee R = 1$. Now,

$$R \subseteq \bigcup_{i \neq j} \pi_i^{-1}(p_i^{\perp i}) \cup \pi_j^{-1}(e(p_j)^c \vee b_j).$$

Note that for all $a \in \mathcal{L}_i$ with $a \subseteq e(p_j)$, we have $e(p_j) \cap (a \vee e(p_j)^{\perp j}) = a$ (see Corollary 6.3). Therefore, $a \vee e(p_j)^{\perp j} = a \cup e(p_j)^{\perp j}$. Hence, we find that

$$R \subseteq \bigcup_{i \neq j} \pi_i^{-1}(p_i^{\perp i}) \cup \pi_j^{-1}(e(p_j)^c \cup b_j).$$

As a consequence, since $\vee R = 1$, from Axiom P2 we find that $b_j = e(p_j)$.

(4) Let $p \in \Sigma$ and $s \in \prod_{i=1}^n e(p_i)$. By part 3, $\phi(p[e(p_1)]) = q^1[e(p_1)]$. Therefore, there is $r_1 \in e(p_1)$ such that $\phi(p[r_1])_1 = s_1$. Let $k \leq n$ and $r_1 \in e(p_1), \dots, r_k \in e(p_k)$ such that $\phi(p[r_1, \dots, r_k])_i = s_i$, for all $i \leq k$, and such that $\phi(p[r_1, \dots, r_k])_{k+1}$ is different from s_{k+1} . By part 3,

$$\phi(p[r_1, \dots, r_k, e(p_{k+1})]) = q^{k+1}[s_1, \dots, s_k, e(p_{k+1})].$$

Hence there is $r_{k+1} \in e(p_{k+1})$ such that $\phi(p[r_1, \dots, r_{k+1}])_i = s_i$, for all i between 1 and $k+1$. As a consequence, from Lemma 6.7, ϕ is surjective. $\square$

9. COVERING PROPERTY

In this Section we prove that the top element $\bigvee_\alpha \mathcal{L}_\alpha$ has the covering property if and only at most one $\mathcal{L}_\alpha$ is not a power set. We reproduces the analogue result concerning the bottom element $\bigwedge_\alpha \mathcal{L}_\alpha$ which is due to Aerts [1]. Moreover, for $\mathcal{L}_i = \text{MO}_{\Sigma_i}$ ($i = 1, 2$) and $T = \text{Aut}(\mathcal{L}_1) \times \text{Aut}(\mathcal{L}_2)$, we prove that there is only one $\mathcal{L} \in \mathcal{S}_T(\mathcal{L}_1, \mathcal{L}_2)$ with the covering property.

Theorem 9.1 (D. Aerts, [1]). *Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$ with for all $\alpha \in \Omega$, $\mathcal{L}_\alpha$ orthocomplemented. If $\bigwedge_\alpha \mathcal{L}_\alpha$ has the covering property or is orthomodular, then there is at most one $\beta \in \Omega$ such that $\mathcal{L}_\beta \neq 2^{\Sigma_\beta}$.*

Proof. Let $\mathcal{L} \in \mathbf{Cl}(\Sigma)$ be orthocomplemented and let $p, q \in \Sigma$ such that $p \vee q = p \cup q$. Define $x := q^\perp \cap (p \vee q)$, then $0 \subseteq x \subseteq p$. If $\mathcal{L}$ has the covering property, then $1 \neq x^\perp$, whereas if $\mathcal{L}$ is orthomodular, $x \vee q = p \vee q$. As a consequence, $x = p$, hence $p \perp q$.

Let $\beta \in \Omega$. Suppose that $\mathcal{L}_\beta \neq 2^{\Sigma_\beta}$. Then there are two non orthogonal atoms, say r_β and s_β . Let $r_\gamma, s_\gamma \in \Sigma_\gamma$ for some $\gamma \in \Omega$ different from β , and let $p, q \in \prod_\alpha \Sigma_\alpha$ defined as $p_\alpha = q_\alpha$, for all $\alpha \neq \beta, \gamma$, and $p_\alpha = r_\alpha$ and $q_\alpha = s_\alpha$ if $\alpha = \beta$ or γ . By Lemma 7.4, $p \vee q = p \cup q$. Therefore, by what precedes, since $\#$ is an orthocomplementation of $\bigodot_\alpha \mathcal{L}_\alpha$, $p \# q$, hence by Definition 3.1, $r_\gamma \perp_\gamma s_\gamma$. As a consequence, $\mathcal{L}_\gamma = 2^{\Sigma_\gamma}$. $\square$

Proposition 9.2. *Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$. If $\bigodot_\alpha \mathcal{L}_\alpha$ has the covering property, then all $\mathcal{L}_\alpha$'s have the covering property.*

Proof. Let $a_\beta \in \mathcal{L}_\beta$, $q_\beta \in \Sigma_\beta$ not in a_β , and $p \in \prod_\alpha \Sigma_\alpha$. By the covering property, we find that $p[q_\beta] \vee p[a_\beta] \succ p[a_\beta]$; Whence by Axiom P3, $q_\beta \vee a_\beta \succ a_\beta$. $\square$

Remark 9.3. In the next theorem, we assume that each $\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)$ with $\mathcal{L}_\alpha \neq 2^{\Sigma_\alpha}$ contains $\mathbf{MO}_4$. This is for instance the case if $\mathcal{L}_\alpha$ is orthocomplemented orthomodular with the covering property. Indeed, if $\mathcal{L}_\alpha \neq 2^{\Sigma_\alpha}$, there is an atom p_α which is not central, hence such that $e(p_\alpha)$ contains at least two atoms, say r and s . Moreover, $r \vee s$ contains at least three atoms, for $\mathcal{L}_\alpha$ has the covering property. Finally, since $\mathcal{L}_\alpha$ is orthomodular, $[0, r \vee s]$ is orthocomplemented, hence contains at least four atoms.

Theorem 9.4. *Let $\{\mathcal{L}_\alpha \in \mathbf{Cl}(\Sigma_\alpha)\}_{\alpha \in \Omega}$.*

- (1) *If each $\mathcal{L}_\alpha$ has the covering property and there is at most one $\beta \in \Omega$ such that $\mathcal{L}_\beta \neq 2^{\Sigma_\beta}$, then $\bigodot_\alpha \mathcal{L}_\alpha$ has the covering property.*
- (2) *Suppose that each $\mathcal{L}_\alpha$ different from 2^{Σ_α} contains $\mathbf{MO}_4$. If $\bigodot_\alpha \mathcal{L}_\alpha$ has the covering property, then there is at most one $\beta \in \Omega$ such that $\mathcal{L}_\beta \neq 2^{\Sigma_\beta}$.*

Proof. (1) Let $\Sigma = \prod_\alpha \Sigma_\alpha$, $a \in \bigodot_\alpha \mathcal{L}_\alpha$, $q \in \Sigma$ not in a , and $R = q \cup a$. By Lemma 5.1,

$$q \vee_{\bigodot} a = \vee_\beta R = \bigcup_{p \in \Sigma} p[\vee R_\beta[p]].$$

Now, by definition,

$$R_\beta[p] = \pi_\beta(p[\Sigma_\beta] \cap (q \cup a)) = q_\beta[p] \cup a_\beta[p],$$

and $q_\beta[p] = q_\beta$ if $p \in q[\Sigma_\beta]$, and $q_\beta[p] = \emptyset$ otherwise. Hence, for all p not in $q[\Sigma_\beta]$, $R_\beta[p] = a_\beta[p] \in \mathcal{L}_\beta$, and if $p \in q[\Sigma_\beta]$, then $R_\beta[p] = q_\beta \cup a_\beta[p]$. Therefore, we find that

$$q \cup a \subseteq q \vee_{\bigodot} a = q[q_\beta \vee a_\beta[q]] \bigcup_{p \notin q[\Sigma_\beta]} p[a_\beta[p]] \subseteq a \cup q[q_\beta \vee a_\beta[q]].$$

As a consequence, $q \vee_{\bigodot} a \succ a$.

(2) Let $\beta \neq \gamma \in \Omega$. Suppose that neither $\mathcal{L}_\beta = 2^{\Sigma_\beta}$ nor $\mathcal{L}_\gamma = 2^{\Sigma_\gamma}$. Let $p, q, r, s, t \in \Sigma$ such that $p_\alpha = q_\alpha = r_\alpha = s_\alpha = t_\alpha$, for all α different from β and γ , and such that $t_\beta = p_\beta$ and $t_\gamma = q_\gamma$. Assume moreover that for $\alpha = \beta$ and for $\alpha = \gamma$, $p_\alpha, q_\alpha, r_\alpha$ and s_α are all different and that $p_\alpha \vee q_\alpha$ covers $p_\alpha, q_\alpha, r_\alpha$ and s_α . By Definition 2.11, $a = \{p, q, r\}$ and $b = \{p, q, r, s\}$ are in $\bigodot_\alpha \mathcal{L}_\alpha$. Let $R^0 = a \cup t$. Then (see Lemma 5.1),

$$\begin{aligned} R^1 &:= \vee_\gamma R^0 = R^0 \cup p[p_\gamma \vee q_\gamma] \\ R^2 &:= \vee_\beta R^1 = R^0 \cup p[p_\gamma \vee q_\gamma] \cup q[p_\beta \vee q_\beta] \cup r[p_\beta \vee q_\beta] \cup s[p_\beta \vee q_\beta] \\ R^3 &:= \vee_\gamma R^2 = p[p_\beta \vee q_\beta, p_\gamma \vee q_\gamma]. \end{aligned}$$

Hence, by Lemma 5.1, $a \vee_{\mathcal{O}} t = R^3 \supsetneq b \supsetneq a$, therefore $\bigvee_{\alpha} \mathcal{L}_{\alpha}$ has not the covering property. $\square$

Definition 9.5. Let Σ_1, Σ_2 be sets, $\mathcal{L}_1 = \text{MO}_{\Sigma_1}$, and $\mathcal{L}_2 = \text{MO}_{\Sigma_2}$ (see Definition 5.3). Then,

$$\mathcal{L}_1 \circ \mathcal{L}_2 := \mathcal{L}_1 \bigotimes \mathcal{L}_2 \cup \{R \in \Xi(\Sigma_1 \times \Sigma_2); |R| = 3\}.$$

Theorem 9.6. Let Σ_1 and Σ_2 be sets, $\mathcal{L}_1 = \text{MO}_{\Sigma_1}$, $\mathcal{L}_2 = \text{MO}_{\Sigma_2}$, $T = \text{Aut}(\mathcal{L}_1) \times \text{Aut}(\mathcal{L}_2)$, and $\mathcal{L} \in \mathcal{S}_T(\mathcal{L}_1, \mathcal{L}_2)$. Then $\mathcal{L}$ has the covering property $\Leftrightarrow \mathcal{L} = \mathcal{L}_1 \circ \mathcal{L}_2$.

Proof. ($\Leftarrow$) Let $\Sigma = \Sigma_1 \times \Sigma_2$, $\Xi = \Xi(\Sigma_1 \times \Sigma_2)$, and $a \in \mathcal{L}_1 \circ \mathcal{L}_2$. Then $a \in \Xi$ and $|a| = 3$, or $a \in \mathcal{L}_1 \bigotimes \mathcal{L}_2$. Hence one of the following cases holds.

- (1) $a \in \Sigma$.
- (2) $a \in \Xi$ and $|a| = 2$ or 3 .
- (3) $a = p_1 \times \Sigma_2$ or $a = \Sigma_2 \times p_2$ for some $p = (p_1, p_2) \in \Sigma$.
- (4) $a = p_1 \times \Sigma_2 \cup \Sigma_1 \times p_2$, for some $p \in \Sigma$ (i.e. a is a coatom).

Hence, obviously $\mathcal{L}_1 \circ \mathcal{L}_2$ has the covering property.

($\Rightarrow$) Let $\mathcal{L} \in \mathcal{S}(\mathcal{L}_1, \mathcal{L}_2)$ with the covering property, and $R \subseteq \Sigma$.

(1) By Lemma 5.1 and Axiom P2, if $R \notin \Xi$ (i.e. there are $p^1, p^2 \in R$ with $p_i^1 = p_i^2$ for $i = 1$ or 2), then $\vee R \in \mathcal{L}_1 \bigotimes \mathcal{L}_2$.

(2) Suppose now that $R \in \Xi$. By Lemma 7.4 part 1, if $|R| \leq 2$, then $R \in \mathcal{L}_1 \bigotimes \mathcal{L}_2$, hence $R \in \mathcal{L}$. Moreover, if $|R| \geq 3$, then for all $s \in \Sigma$ not in R with $s_1 \in \pi_1(R)$ or $s_2 \in \pi_2(R)$, we have $\vee(R \cup s) = 1$.

(3) Suppose that $R \in \Xi$ and that $|R| = 3$. Write $a := \vee R$. **Claim:** $a \neq 1$ [*Proof:* Write $R = \{p, q, r\}$ and suppose that $a = 1$. As we have seen $R^0 := \{p, q\} \in \mathcal{L}$. Hence

$$1 = a = r \vee R^0 \supsetneq p_1 \times \Sigma_2 \cup \Sigma_1 \times q_2 \supsetneq R^0,$$

a contradiction since $\mathcal{L}$ has the covering property.]

As a consequence, from part 2, $1 \succ a$. Moreover, $a \in \Xi$. We write a as $a = \{p_1 \times f(p_1); p_1 \in \pi_1(a)\}$. Hence, f is injective.

(4) **Claim:** $a = R$. [*Proof:* First, note that any bijection of Σ_2 induces an automorphism of $\mathcal{L}_2$. Suppose that $a \neq R$, hence $|a| \geq 4$. Let $v_2 \in \text{Aut}(\mathcal{L}_2)$ such that its restriction to $\pi_2(a)$ is different from the identity, and with at least three fixed points in $\pi_2(a)$. By Axiom P4, there is $u \in \text{Aut}(\mathcal{L})$ such that on Σ , u equals $\text{id} \times v_2$. Hence $c := \{(p_1, v_2 \circ f(p_1)); p_1 \in \pi_1(a)\} \in \mathcal{L}$ and $c \cap a \neq a$, therefore 1 does not cover $c \cap a$. Moreover $|c \cap a| \geq 3$; whence a contradiction by part 3.]

(5) Finally, suppose that $R \in \Xi$ and $|R| \geq 4$. Let $b = \vee R$. **Claim:** $b = 1$. [*proof:* Let $R_0 \in \Xi$ with $R_0 \subseteq R$ and $|R_0| = 3$. By what precedes, $R_0 \in \mathcal{L}$ and $1 \succ R_0$. As a consequence, $b = 1$.] $\square$

10. AN EXAMPLE WITH THE COVERING PROPERTY

Let $\mathcal{H}_1$ and $\mathcal{H}_2$ be complex Hilbert spaces and $T = \text{U}(\mathcal{H}_1) \times \text{U}(\mathcal{H}_2)$. In this section, we give an example in $\mathcal{S}_T(\text{P}(\mathcal{H}_1), \text{P}(\mathcal{H}_2))$, denoted by $\text{P}(\mathcal{H}_1) \oplus \text{P}(\mathcal{H}_2)$, which has the covering property. Moreover, $\text{P}(\mathcal{H}_1) \oplus \text{P}(\mathcal{H}_2)$ is coatomistic, but, as expected, the dual has not the covering property, i.e. $\text{P}(\mathcal{H}_1) \oplus \text{P}(\mathcal{H}_2)$ is not a DAC-lattice.

Definition 10.1. Let $\mathcal{H}_1$ and $\mathcal{H}_2$ be complex Hilbert spaces, $\Sigma = \Sigma_{\mathcal{H}_1} \times \Sigma_{\mathcal{H}_2}$, $\mathcal{L}_1 = \text{P}(\mathcal{H}_1)$, $\mathcal{L}_2 = \text{P}(\mathcal{H}_2)$, and $\Sigma_{\otimes}$ the set of one-dimensional subspaces of $\mathcal{H}_1 \otimes \mathcal{H}_2$.

For $V \in \mathbf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$, define $\Sigma_{\Downarrow}[V] := \{(p_1, p_2) \in \Sigma; p_1 \otimes p_2 \in V\}$. Then,

$$\mathcal{L}_1 \boxplus \mathcal{L}_2 := \{\Sigma_{\Downarrow}[V]; V \in \mathbf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)\},$$

ordered by set-inclusion. For $A \subseteq \Sigma$, we write $A^\perp := \{q \in \Sigma_{\otimes}; \langle q, p_1 \otimes p_2 \rangle = 0, \forall (p_1, p_2) \in A\}$, where $\langle - | - \rangle$ denotes the scalar product in $\mathcal{H}_1 \otimes \mathcal{H}_2$. Moreover, we denote the set of antilinear maps from $\mathcal{H}_1$ to $\mathcal{H}_2$ by $\mathcal{A}(\mathcal{H}_1, \mathcal{H}_2)$.

Proposition 10.2. *Let m and n be integers, $\mathcal{H}_1 = \mathbb{C}^m$, $\mathcal{H}_2 = \mathbb{C}^n$, $\mathcal{L}_1 = \mathbf{P}(\mathcal{H}_1)$, $\mathcal{L}_2 = \mathbf{P}(\mathcal{H}_2)$, and $\Sigma = \Sigma_{\mathcal{H}_1} \times \Sigma_{\mathcal{H}_2}$. For $A \in \mathcal{A}(\mathcal{H}_1, \mathcal{H}_2)$, define $X_A \subseteq \Sigma$ as $X_A := \cup\{p_1 \times (A(p_1)^{\perp_2}); p_1 \in \Sigma_1\}$. Then,*

$$\mathcal{L}_1 \boxplus \mathcal{L}_2 = \{\cap \omega; \omega \subseteq \{X_A; A \in \mathcal{A}(\mathcal{H}_1, \mathcal{H}_2)\}\}.$$

Proof. Let $\{e_i^1\}_{1 \leq i \leq m}$ and $\{e_j^2\}_{1 \leq j \leq n}$ denote the canonical basis of $\mathbb{C}^m$ and $\mathbb{C}^n$ respectively.

Let $v \in \mathbb{C}^m \otimes \mathbb{C}^n$ and $p = (p_1, p_2) \in \Sigma$ with $p_1 = \mathbb{C}w_1$ and $p_2 = \mathbb{C}w_2$. Write v , w_1 and w_2 as

$$v = \sum_{i=1}^m \sum_{j=1}^n s_{ij} e_i^1 \otimes e_j^2, \quad w_1 = \sum_{i=1}^m \lambda_i e_i^1, \quad \text{and} \quad w_2 = \sum_{j=1}^n \mu_j e_j^2.$$

Let

$$\lambda = (\lambda_1, \dots, \lambda_m)^\top \text{ and } \mu = (\mu_1, \dots, \mu_n)^\top.$$

Let S be the $m \times n$ matrix defined as $S_{ij} = s_{ij}$. Then $p \in \Sigma_{\Downarrow}[v^\perp]$ if and only if $\langle w_1 \otimes w_2, v \rangle = 0$, hence if and only if $\bar{\mu}^\top S^\top \bar{\lambda} = 0$. Let A be the antilinear map defined by the matrix $S^\top$. Then, $p \in \Sigma_{\Downarrow}[v^\perp]$ if and only if $p_2 \in A(p_1)^\perp$, that is if and only if $p \in X_A$. As a consequence, $\Sigma_{\Downarrow}[v^\perp] = X_A$.

On the other hand, if $A \in \mathcal{A}(\mathcal{H}_1, \mathcal{H}_2)$, then $X_A = \Sigma_{\Downarrow}[v^\perp]$, where v is given by the formula above with $s_{ij} = (A^\top)_{ij}$. $\square$

Remark 10.3. Let $\mathcal{H}$ be a complex Hilbert space of dimension ≥ 3 . Then, by Wigner's theorem (see [5], Theorem 14.3.6), any ortho-automorphism of $\mathbf{P}(\mathcal{H})$ is induced by a unitary or antiunitary map on $\mathcal{H}$. Note that if v_1 is a unitary map on $\mathcal{H}_1$ and v_2 is an antiunitary map on $\mathcal{H}_2$, then $v = v_1 \times v_2$ does not induce an automorphism of $\mathcal{L}_1 \boxplus \mathcal{L}_2$. Indeed, let X_A be a coatom. Then

$$v(X_A) = \bigcup_{p_1 \in \Sigma_1} v_1(p_1) \times v_2(A(p_1)^{\perp_2}) = \bigcup_{p_1 \in \Sigma_1} p_1 \times ((v_2 \circ A \circ v_1^{-1}(p_1))^{\perp_2}).$$

Now, since A and v_2 are antilinear and v_1 is linear, it follows that $v_2 \circ A \circ v_1^{-1}$ is linear, hence $v(X_A)$ is not a coatom of $\mathcal{L}_1 \boxplus \mathcal{L}_2$.

Theorem 10.4. *Let $\mathcal{H}_1$ and $\mathcal{H}_2$ be complex Hilbert spaces, $\Sigma = \Sigma_{\mathcal{H}_1} \times \Sigma_{\mathcal{H}_2}$, $\mathcal{L}_1 = \mathbf{P}(\mathcal{H}_1)$, $\mathcal{L}_2 = \mathbf{P}(\mathcal{H}_2)$, and $T = \mathbf{U}(\mathcal{H}_1) \times \mathbf{U}(\mathcal{H}_2)$. Then*

- (1) *for all $A \subseteq \Sigma$, we have $\vee_{\boxplus} A = \Sigma_{\Downarrow}[A^{\perp\perp}]$,*
- (2) $\mathcal{L}_1 \boxplus \mathcal{L}_2 \in \mathcal{S}_T(\mathcal{L}_1, \mathcal{L}_2)$,
- (3) $\mathcal{L}_1 \boxplus \mathcal{L}_2$ *has the covering property and is coatomistic, but, if $\mathcal{L}_1 \neq 2 \neq \mathcal{L}_2$ (i.e. the dimension of $\mathcal{H}_1$ and $\mathcal{H}_2$ is ≥ 2), the dual has not the covering property,*
- (4) $\mathcal{L}_1 \boxtimes \mathcal{L}_2 = \{\Sigma_{\Downarrow}[V]; V \in \mathbf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2), V = \Sigma_{\Downarrow}[V]^{\perp\perp}, V^\perp = \Sigma_{\Downarrow}[V^\perp]^{\perp\perp}\}$ *(i.e. both V and $V^\perp$ are spanned by product vectors),*
- (5) *if $\mathcal{L}_1 \neq 2 \neq \mathcal{L}_2$, then $\mathcal{L}_1 \boxtimes \mathcal{L}_2 \subsetneq \mathcal{L}_1 \boxplus \mathcal{L}_2 \subsetneq \mathcal{L}_1 \boxdot \mathcal{L}_2$.*

Proof. (1) Follows directly from Definition 10.1.

(2) Obviously, $\mathcal{L}_1 \oplus \mathcal{L}_2 \in \mathbf{Cl}(\Sigma)$. Let $a \in \mathcal{L}_1 \otimes \mathcal{L}_2$. By Definition 3.1,

$$a^\# = \Sigma_\downarrow[a^\perp].$$

Hence, $a^\# \subseteq a^\perp$, $a^{\perp\perp} \subseteq a^{\#\perp}$, thus $\Sigma_\downarrow[a^{\perp\perp}] \subseteq a^{\#\#} = a$, therefore $a = \Sigma_\downarrow[a^{\perp\perp}]$. As a consequence, $\mathcal{L}_1 \otimes \mathcal{L}_2 \subseteq \mathcal{L}_1 \oplus \mathcal{L}_2$.

Let $V \in \mathbf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$ such that $\Sigma_\downarrow[V] = p_1 \times A_2$. Then

$$p_1 \otimes (A_2^{\perp_2 \perp_2}) = (p_1 \otimes A_2)^{\perp\perp} \subseteq V,$$

therefore $A_2^{\perp_2 \perp_2} \subseteq A_2$, hence $A_2 \in \mathcal{L}_2$. As a consequence, Axiom P3 holds.

Axiom P4 with $T = \mathbf{U}(\mathcal{H}_1) \times \mathbf{U}(\mathcal{H}_2)$ holds in $\mathbf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$, therefore obviously also in $\mathcal{L}_1 \oplus \mathcal{L}_2$.

(3) The covering property holds in $\mathbf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$ (see [13], Theorem 34.2), hence, by part 1, also in $\mathcal{L}_1 \oplus \mathcal{L}_2$. Moreover, since $\mathbf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$ is coatomistic, so is $\mathcal{L}_1 \oplus \mathcal{L}_2$.

Finally, let $p \in \Sigma$. Then $x = \Sigma_\downarrow[p^\perp] = p^\#$ is coatom of $\mathcal{L}_1 \oplus \mathcal{L}_2$. Now, there is $R \in \Xi(\Sigma)$ (see Definition 5.3) with $|R| = 2$, such that $x \cap R = \emptyset$. By Lemma 7.4, $R \in \mathcal{L}_1 \oplus \mathcal{L}_2$. Moreover $x \vee_\oplus R = 1$ since x is a coatom. Hence, writing $R = \{p, q\}$, and the order relation, meet, join, bottom and top elements in $(\mathcal{L}_1 \oplus \mathcal{L}_2)^*$ by $\leq_*$, $\wedge_*$, $\vee_*$, 0_* , and 1_* respectively, we find that $x \wedge_* R = 0_*$ and $x \vee_* R = 1_* \not\geq_* p \not\geq_* R$. Therefore, $(\mathcal{L}_1 \oplus \mathcal{L}_2)^*$ has not the covering property.

(4) Let $p \in \Sigma$ and $q \in p^{\#\perp}$. Write $q = \mathbb{C}v$ with $v \in \mathcal{H}_1 \otimes \mathcal{H}_2$. For $i = 1$ and $i = 2$, let $\{w_i^k\}$ be an ortho-basis of $p_i^{\perp i}$, and let $x_i \in p_i$ (i.e. $p_i = \mathbb{C}x_i$). Then v can be decomposed as

$$v = \alpha x_1 \otimes x_2 + \sum_{k_2} \beta_{k_2} x_1 \otimes w_2^{k_2} + \sum_{k_1} \beta_{k_1} w_1^{k_1} \otimes x_2 + \sum_{l_1 l_2} \gamma_{l_1 l_2} w_1^{l_1} \otimes w_2^{l_2}.$$

Now, $p^\# = p_1^{\perp 1} \times \Sigma_{\mathcal{H}_2} \cup \Sigma_{\mathcal{H}_1} \times p_2^{\perp 2}$. Hence, since $v \in p^{\#\perp}$, we find that $\beta_{k_1} = \beta_{k_2} = \gamma_{l_1 l_2} = 0$, for all k_1, k_2, l_1 and l_2 . Therefore, $v \in p$, hence $p^{\#\perp} = p$.

Let $a \in \mathcal{L}_1 \otimes \mathcal{L}_2$. From part 1, $a = \Sigma_\downarrow[a^{\perp\perp}]$. On the other hand, $a^\# = \Sigma_\downarrow[a^\perp]$. Now, $a = a^{\#\#} = \cap\{p^\#; p \in a^\#\}$. Hence, by what precedes,

$$a^\perp = (\cup\{p^{\#\perp}; p \in a^\#\})^{\perp\perp} = a^{\#\perp\perp},$$

therefore $a^\perp$ is also spanned by product vectors.

Let $V \in \mathbf{P}(\mathcal{H}_1 \otimes \mathcal{H}_2)$ such that both V and $V^\perp$ are spanned by product vectors. Let $a := \Sigma_\downarrow[V]$ and $b := \Sigma_\downarrow[V^\perp]$. Since $\Sigma_\downarrow[V]^{\perp\perp} = V$ and $\Sigma_\downarrow[V^\perp]^{\perp\perp} = V^\perp$, we find that

$$\Sigma_\downarrow[\Sigma_\downarrow[V]^\perp] = \Sigma_\downarrow[V^\perp] \text{ and } \Sigma_\downarrow[\Sigma_\downarrow[V^\perp]^\perp] = \Sigma_\downarrow[V].$$

Therefore $a^\# = b$ and $b^\# = a$. As a consequence, $a \in \mathcal{L}_1 \otimes \mathcal{L}_2$.

(5) By part 4, $\mathcal{L}_1 \otimes \mathcal{L}_2 \neq \mathcal{L}_1 \oplus \mathcal{L}_2$. On the other hand, by Theorem 9.4, $\mathcal{L}_1 \otimes \mathcal{L}_2$ has not the covering property, whereas by part 3, $\mathcal{L}_1 \oplus \mathcal{L}_2$ has the covering property. As a consequence, $\mathcal{L}_1 \oplus \mathcal{L}_2 \neq \mathcal{L}_1 \otimes \mathcal{L}_2$. $\square$

ACKNOWLEDGMENTS

An important part of this work was done during a stay at McGill. In this connection, I would like to thanks M. Barr for his hospitality.

REFERENCES

- [1] D. Aerts. Description of many separated physical entities without the paradoxes encountered in quantum mechanics. *Found. Phys.*, 12(12):1131–1170, 1982.
- [2] D. Aerts. The missing elements of reality in the description of quantum mechanics of the EPR paradox situation. *Helv. Phys. Acta*, 57(4):421–428, 1984.
- [3] M. Barr. **-autonomous categories*, volume 752 of *Lecture Notes in Mathematics*. Springer, Berlin, 1979. With an appendix by Po Hsiang Chu.
- [4] G. Birkhoff and J. v. Neumann. The logic of quantum mechanics. *Ann. of Math. (2)*, 37(4):823–843, 1936.
- [5] C.-A. Faure and A. Frölicher. *Modern projective geometry*, volume 521 of *Mathematics and its Applications*. Kluwer Academic Publishers, Dordrecht, 2000.
- [6] C.-A. Faure, D. J. Moore, and C. Piron. Deterministic evolutions and Schrödinger flows. *Helv. Phys. Acta*, 68(2):150–157, 1995.
- [7] G. A. Fraser. The semilattice tensor product of distributive lattices. *Trans. Amer. Math. Soc.*, 217:183–194, 1976.
- [8] A. S. Golfin. *Representations and products of lattices*. PhD thesis, U. Mass., Amherst, 1987.
- [9] G. Grätzer and F. Wehrung. A new lattice construction: the box product. *J. Algebra*, 221(1):315–344, 1999.
- [10] B. Ischi. A characterization of the Aerts product of Hilbertian lattices. Submitted, available on <http://arxiv.org/abs/math-ph/0405048>.
- [11] B. Ischi. Orthocomplementation and compound systems. Submitted, available on <http://arxiv.org/abs/quant-ph/0410085>.
- [12] B. Ischi. Endomorphisms of the separated product of lattices. *Internat. J. Theoret. Phys.*, 39(11):2559–2581, 2000.
- [13] F. Maeda and S. Maeda. *Theory of symmetric lattices*. Die Grundlehren der mathematischen Wissenschaften, Band 173. Springer-Verlag, New York, 1970.
- [14] J. v. Neumann. *Mathematical foundations of quantum mechanics*. Princeton University Press, Princeton, 1955. Translated by Robert T. Beyer.
- [15] C. Piron. Axiomatique quantique. *Helv. Phys. Acta*, 37:439–468, 1964.
- [16] C. Piron. *Foundations of quantum physics*. W. A. Benjamin, Inc., Advanced Book Program, Reading, Mass.-London-Amsterdam, 1976. Mathematical Physics Monograph Series, 19.
- [17] S. Pulmannová. Tensor product of quantum logics. *J. Math. Phys.*, 26(1):1–5, 1985.
- [18] G. J. Seal. Cartesian closed topological categories and tensor products. *Appl. Categ. Structures*, 13(1):37–37, 2005.
- [19] Z. Shmueli. The structure of Galois connections. *Pacific J. Math.*, 54(2):209–225, 1974.

BORIS ISCHI, LABORATOIRE DE PHYSIQUE DES SOLIDES, UNIVERSITÉ PARIS-SUD, BÂTIMENT 510,
91405 ORSAY, FRANCE
E-mail address: ischi@kalymnos.unige.ch